

গণপ্রজাতন্ত্রী বাংলাদেশ সরকার
কৃষি মন্ত্রণালয়
মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট
কৃষি খামার সড়ক, ঢাকা-১২১৫।
www.srdi.gov.bd

নম্বর ১২.০৩.০০০০.০৩১.৯৯.০০৩.১৮.৭

তারিখ ২৮ তাদ্র ১৪২৬

১২ সেপ্টেম্বর ২০১৯


বিষয়: ২০১৮-১৯ অর্থ বছরের বার্ষিক কাজের অগ্রগতি প্রতিবেদনের জন্য তথ্য প্রেরণ প্রসংগে।

সূত্র: ১২.০৩.০০০০.০০২.০৬.০০৪.১৯.৬২৮; তারিখ: ০৪-০৯-২০১৯ খ্রি.

উপর্যুক্ত বিষয় ও সূত্রেয় পত্রের প্রেক্ষিতে জানানো যাচ্ছে যে, এসআরডিআই-এর বার্ষিক কর্মসম্পাদন চুক্তির (২০১৯-২০) আবশ্যিক কৌশলগত উদ্দেশ্যসমূহ মোতাবেক, ২০১৮-১৯ অর্থ বছরের বার্ষিক কারিগরি কাজের অগ্রগতি প্রতিবেদন প্রণয়নের বাধ্যবাধকতা রয়েছে। সংযুক্ত ছক (সফট কপি) মোতাবেক দপ্তর/অফিস সমূহের তথ্য উপাত্ত জরুরী প্রয়োজন।

এমতাবস্থায়, প্রেরিত ছক মোতাবেক তথ্য উপাত্ত আগামী ২৫-০৯-২০১৯ খ্রি. তারিখের মধ্যে ড. মো: আলতাফ হোসেন, প্রধান বৈজ্ঞানিক কর্মকর্তা ও আহবায়ক, বার্ষিক প্রতিবেদন প্রণয়ন কমিটি ই-মেইল (altajolly63@gmail.com), মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট-এর নিকট সফট ও হার্ড কপি আকারে প্রেরণের জন্য আদেশ করা হলো।

সংযুক্ত: বর্ণনামতে।



১২-৯-২০১৯
বিধান কুমার ভান্ডার
পরিচালক

সদয় অবগতি ও কার্যার্থে প্রেরণ করা হল:

- ১) মুখ্য বৈজ্ঞানিক কর্মকর্তা, উন্নয়ন বিভাগ/মৃত্তিকা পরীক্ষা বিভাগ, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।
- ২) প্রকল্প পরিচালক, নগর কৃষি উৎপাদন সহায়ক (পাইলট) প্রকল্প, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।
- ৩) প্রকল্প পরিচালক, মৃত্তিকা গবেষণা এবং গবেষণা সুবিধা জোরদারকরণ (এসআরএসআরএফ) প্রকল্প, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।
- ৪) প্রকল্প পরিচালক, গোপালগঞ্জ-খুলনা-বাগেরহাট-সাতক্ষীরা-পিরোজপুর কৃষি উন্নয়ন (এসআরডিআই অংগ) প্রকল্প, খুলনা।
- ৫) কর্মসূচি পরিচালক, ভাম্যমান মৃত্তিকা পরীক্ষাগারের (এমএসটিএল) মাধ্যমে সরেজমিনে কৃষকের মাটি পরীক্ষার ভিত্তিতে সুমম সার সুপারিশ কার্যক্রম জোরদারকরণ কর্মসূচি, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।
- ৬) প্রধান বৈজ্ঞানিক কর্মকর্তা, কেন্দ্রীয় গবেষণাগার/প্রশিক্ষণ শাখা/মৃত্তিকা জরিপ ব্যাখ্যা শাখা/মৃত্তিকা ও ভূমি শ্রেণিবিন্যাস জরিপ শাখা/মৃত্তিকা কোরিলেশন শাখা/উপজেলা নির্দেশিকা সেল, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।
- ৭) প্রধান বৈজ্ঞানিক কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, আঞ্চলিক কার্যালয়, ঢাকা/রাজশাহী/কুমিল্লা/খুলনা/ সিলেট/বরিশাল।
- ৮) প্রধান বৈজ্ঞানিক কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, আঞ্চলিক গবেষণাগার, ঢাকা/রাজশাহী/কুমিল্লা/খুলনা/ ময়মনসিংহ।
- ৯) ইনোভেশন অফিসার, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট (ওয়েব সাইটে প্রকাশের অনুরোধসহ)।
- ১০) প্রধান বৈজ্ঞানিক কর্মকর্তা/ভারপ্রাপ্ত কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, মৃত্তিকা সংরক্ষণ ও পানি বিভাজিকা ব্যবস্থাপনা কেন্দ্র,

মান্দরবান/লবণাক্ততা ব্যবস্থাপনা ও গবেষণা কেন্দ্র, বটিয়াঘাটা, খুলনা।

১১) উর্ধ্বতন বৈজ্ঞানিক কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, জেলা কার্যালয়, ঢাকা/ময়মনসিংহ/জামালপুর/

টাংগাইল/ফরিদপুর/রাজশাহী/পাবনা/বগুড়া/রংপুর/দিনাজপুর/কুমিল্লা/চট্টগ্রাম/রাঙ্গামাটি/নোয়াখালী/খুলনা/যশোর/কুষ্টিয়া/পটুয়াখালী/সিলেট/মৌলভীবাজার।

১২) ভারপ্রাপ্ত কর্মকর্তা, ডাটা প্রসেসিং ও পরিসংখ্যান শাখা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।

১৩) জনাব সৈয়দ আহসান রেজা চৌধুরী, পরিচালক মহোদয়ের সংযুক্ত কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।

১৪) উর্ধ্বতন মানচিত্রাংকনবিদ, মানচিত্রাংকন শাখা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।

১৫) ভারপ্রাপ্ত কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, আঞ্চলিক গবেষণাগার, জামালপুর/ফরিদপুর/চট্টগ্রাম/সিলেট/
নোয়াখালী/বগুড়া/দিনাজপুর/বরিশাল/কুষ্টিয়া/ঝিনাইদহ।


১৬) পাবলিকেশন এন্ড লিয়ার্জো অফিসার, প্রকাশনা ও রেকর্ড শাখা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।

১৭) সহকারী পরিচালক/প্রশাসনিক কর্মকর্তা/স্টোর অফিসার/সহকারী হিসাবরক্ষণ কর্মকর্তা, মৃত্তিকা সম্পদ উন্নয়ন ইনস্টিটিউট, ঢাকা।

১৮) অফিস নথি।



ANNUAL REPORT 2017-2018



Jaldi series
Bandarban



Slash and Burn for Jhum in CHT



Rabi crops in Ghior



SOIL RESOURCE DEVELOPMENT INSTITUTE
MINISTRY OF AGRICULTURE
MRITTIKA BHABAN, FARMGATE, DHAKA



Government of the People's Republic of Bangladesh

ANNUAL REPORT 2017-2018



SOIL RESOURCE DEVELOPMENT INSTITUTE

MINISTRY OF AGRICULTURE

MRITTIKA BHABAN, FARMGATE, DHAKA-1215

2018

ANNUAL REPORT

2017-2018

Compiled by

Md. Kamaruzzaman
Md. Liaquat Hossain
Dr. Md. Taiabur Rahman
Dr. Md. Altaf Hossain
Dr. Md. Mohiuddin Ahmed
Md. SabbirHossain
Sachindranath Biswas
Monfique Ahmed Chowdhury
Md. Jalaluddin
Dr. Begum Samia Sultana
Dr. Md. Afsar Ali
A.F.M. Monjurul Huq
Principal Scientific Officer
Ameer Md. Zahid
Md. Babul Hossain
Senior Scientific Officer

Sharmin Akter
Scientific Officer

Edited by

Bidhan Kumar Bhandar
Director

Foreword

Annual Report 2017-2018 reflects all the activities of Soil Resource Development Institute (SRDI) for the period of July 2017 to June 2018. Activities include both new programs and continuation of on-going programs, which are presented in simple and concise form for easy understanding of the users.

SRDI has the mandate to make land and soil resources inventory to make location specific database, land use planning, monitoring of soil fertility and salinity status, develop appropriate ameliorative measures for saline and other form of degraded soils, advice farmers on crop suitability, soil test and crop demand based balanced fertilizer use, and management and conservation of hill soils. SRDI also investigates soil physical, chemical, biological and biochemical processes and thereby develop primary soil resource database through soil survey, remote sensing and GIS technique.

SRDI has prepared Soil and Land Resource Utilization Guide (popularly known as UpazilaNirdeshika) for each Upazila of the country which is a unique database for local level agricultural development planning and a unique tool for local level extension workers. Subsequently, SRDI is preparing Union Sahayika on land type wise crops and fertilizer dose to be used at Union level. SRDI has innovated Online Fertilizer Recommendation System through which farmers' from any part of the country can know fertilizer requirement for different crops they want to cultivate. Saline soil management and sloping land management techniques have also been developed by SRDI.

In addition, this Institute provides support services to other NARS institutes, GO and NGO's by providing data, information, maps and reports on soil and crop loss by natural disasters like flood, drought, cyclone etc by using GIS technique.

I hope the findings and explanations presented under different heads in this report will be of great use to all concerned agencies and beneficiaries working in the field of agriculture.

I would like to extend my heartiest thanks to the officers involved in implementing the annual program for the year 2017-2018 with sincere efforts and to those who worked hard to make this report. Any suggestions or recommendation to develop the report would be highly appreciated.

(Bidhan Kumar Bhandar)
Director

Contents

Foreword	iii
Acronyms	iv
Executive Summary	1

Chapter 1: SRDI at a glance

1.0 Brief History

1.1 Mandates of SRDI

1.2 Organogram

Chapter 2: Activities of Different Sections of Head Quarter

2.1 Soil Survey and Land Classification Section

2.2 Soil Survey Interpretation Section

2.3 Correlation Section

2.4 Training Section

2.5 Data Processing and Statistical Section

2.6 Cartography Section

2.7 Central Laboratory

2.8 Upazila Nirdeshika Cell

Chapter 3: Activities of Field Offices

3.1 Updating Upazila Nirdeshika through Semi-detailed Soil Survey

3.2 Union Land, Soil and Fertilizer Recommendation Guide Preparation

3.3 Soil and Water Salinity Monitoring

3.4 Monitoring & Evaluation of Farmers' Service through Mobile Soil Testing Laboratories

3.5 Distribution and Monitoring of Fertilizer Recommendation Card (FRC) based on
Upazila Nirdeshika fertility status

3.6 Technology transfer through demonstration

Chapter 4. Activities of Research Centers

4.1 Soil Conservation and Watershed Management Center (SCWMC)

4.2 Salinity Management and Research Center (SMRC)

Chapter 5. Activities of Regional Laboratories

Chapter 6. Project Activities

Executive Summary

During 2017-18 fiscal year “Upazila Nirdeshika Updating Survey” was carried out at 50 Upazilas where remarkable changes in land use were observed in each Upazila (sub-district). Changes in land types were also found in some cases. It is observed that vegetable cultivation is gaining popularity among the farmers. In this year 30 Upazila Nirdeshika and 90 Union Sahayika (guides) were published. In addition, Technology Transfer activities like fertilizer recommendation cards (FRC) distribution, establishment of demonstration plots through collaboration of static laboratories, mobile soil testing laboratories (MSTL) and district offices resulted substantial yield increase in comparison to unattended farmers plots. In 2017-18, soil analytical service through MSTL was provided to 112 Upazila of the country and provided 5,600 farmers with FRC. MSTL activities needs to be extended to Union level to provide services at grass root level.

Changes in soil fertility due to land use and management practices were observed in monitoring sites. In general, soils are deficient in organic matter and nitrogen. Changes in Phosphorus, Potassium, Calcium, Magnesium, Manganese, Sulfur and Boron were erratic. There is also, evidence of lower pH value in many Upazilas.

In Khulna region, soil salinity in shrimp cultivation area gradually increased from 1990. This salinization may be due to the effect of saline water flooding for long period, slow permeability, presence of highly saline ground water at shallower depth (<1.0m) almost throughout the year and lack of flashing facility after shrimp harvest etc. Water salinity of western side river is more than that of the eastern side. Water salinity of the same stream is less in upper stream and higher in the downstream. Most of the river water remains saline throughout the year and not suitable for irrigation. Few of the river water remain non-saline from the month of August to October.

Soil and water salinity is the major constraint for land use development in Barisal region like other coastal areas of the country. Salinity build up in soil and water of this area is due to the influence of sea water. It is quite natural and a common feature of coastal environment. We cannot prevent this process rather we have to adapt to it. So, management of salinity is the only option for agricultural development here. Monitoring activities need to be widened and strengthened to generate database for setting future strategies to encounter the upcoming challenge of sea level rise.

In Chittagong region, salinity commences with the beginning of dryness of soil (usually, November to December) and reached its peak during the month of February to April and starts to decline with the advance of rainfall and the minimum EC: dS/m reaches during July to September. The variability of soil salinity is fully regulated by rainfall, its intensity, duration, and drainage condition. From the results it is found that the EC value is decreasing over the years. In the coastal areas, lands, which are not yet protected by embankment but flooded regularly, should be protected. Increasing shrimp cultivation may increase the salinity and hamper the crop production in neighboring areas, so it should be cultivated in a planned way. For the cultivation

of rabi crops drainage system should be well. Use of chemical fertilizers based on soil analysis and drainage improvement are needed to increase the cropping intensity and achieve higher yields. Use of salt tolerant varieties should be increased. Re-excavation of small khals (creeks) as rain water reservoirs is needed to use as irrigation water source. Land should not be kept fallow. It should be covered by any crop or straw within suitable cropping system. Bare lands should be ploughed to reduce the evaporation.

Some innovative technologies for slopping hill soil management were generated by Soil Conservation and Watershed Management Centre (SCWMC), Meghla, Bandarban of which Quesungual Slash and Mulch Agroforestry (QSMAS), Bench Terrace for year round crop production, Gabion Check Dam for gully erosion control, Jute Geo-Textile for rehabilitating degraded land, establishment of Hedge Rows in farmer's field etc. for soil erosion control and organic matter conservation were most important ones.

Some innovative technologies for saline soil management was generated by Salinity Management and Research Centre (SMRC), Batiaghata, Khulna of which Pitcher Irrigation with split application of fertilizers in soluble form, Double Mulching and Raised Bed for vegetable cultivation were proved worthy. These techniques can be disseminated to other saline areas.

During 2017-18, static laboratories conducted soil analysis for both physical and chemical parameters, plant and water analysis for chemical parameters and fertilizer samples analysis under different programs. In static laboratories (Central and Regional Labs), a total of 25,842 samples (21,045 soil samples, 261 water samples, 87 plant samples and 4,449 fertilizer samples) were analyzed.

Chapter 1: SRDI at a glance

1. Brief Introduction of Soil Resource Development Institute (SRDI):

The Soil Resource Development Institute (SRDI), an attached department to Ministry of Agriculture originated in 1961 as the East Wing Directorate of the Soil Survey Project of Pakistan under Ministry of Agriculture and Works of the then Government of Pakistan. After emergence of Bangladesh, the then east wing office of the Central Soil Resource Institute started functioning as Department of Soil Survey under Ministry of Agriculture and Forest, Government of the People's Republic of Bangladesh. In 1983, Soil Resource Development Institute (SRDI) was established under the Ministry of Agriculture and Forest by reorganizing the then Department of Soil Survey. From 1986 onwards through successful completion of several projects the activities of SRDI has increased manifold. It is now prepared to face the challenges of future to make breakthrough in crop production through improved soil and nutrient management in Bangladesh.

Vision of SRDI:

The vision of SRDI is to ensure judicious and profitable use of scarce land and soil resources of the country and to protect the soil health.

Mission of SRDI:

The missions of SRDI are

- to inventory soil and land resources of the country,
- to classify them according to their potentiality,
- to generate a detail database on soil, land, water and agro-climate,
- to prepare guidelines (Nirdeshika, Shahayika and pustica) for the best uses of the natural resources,
- to identify and manage problem soils and finally
- to ensure active role in sustainable increase of crop production.

1.1. Functions of SRDI:

The functions of the Soil Resource Development Institute are as follows:

- a) Reconnaissance Soil Survey of the whole country on the basis of aerial photo interpretation and field and laboratory investigation of soils;
- b) Detailed/Semi-detailed soil surveys of development project areas and research farms for various beneficiary agencies;
- c) Soil surveys to evaluate irrigation command areas and cropping potentials;
- d) Soil surveys for locating areas of problem soils (e.g., toxic, saline, alkaline or peat soils), soil degradation and erosion (in watershed region) for planning reclamation or watershed management;
- e) Correlation of soils conducted under various surveys;
- f) Chemical analysis of soil, water and plant samples to verify and clarify the field observation;
- g) Analysis of chemical and organic fertilizers to ensure the quality of fertilizers as backup for policy makers;
- h) Interpretation of aerial photos, land sat imageries and topographic maps for soil and land use surveys;
- i) Preparation of various maps and reports on the above-mentioned surveys for publication;
- j) Services to the development agencies by providing basic data on soils, land capability and crop suitability for preparation of both short and long-term agricultural development plans;
- k) Coordination with the beneficiary agencies at local, regional or national levels regarding planning and execution of land use development programs;
- l) Guides on soils and agricultural development possibilities for each Upazila for agricultural extension and research workers;
- m) Provision of soil data for planning irrigation, drainage and reclamation projects;
- n) Selection of suitable sites for specific research/development activities;
- o) Imparting in-service training to the newly recruited technical officers on soil survey, land use planning, cropping potential, etc. and refreshers training to keep the technical officers of the department apprised and acquainted with the up-to date knowledge;
- p) Training of agricultural extension and research workers of various levels on proper utilization of soil survey information. Imparting basic training on various aspects of soils to the students of the agricultural institutions. (Source: Gazette Notification, October, 1983);
- q) Render services to farmers and others by analyzing soil, plant, water and fertilizer samples and recommend location specific fertilizer doses on the basis of soil testing and crop requirements;
- r) Provide assistance in regular monitoring of soil fertility and land productivity activities throughout the country;

- s) Study in soil moisture characteristics to ascertain irrigation needs of different crops;
- t) Launch a regular program for the training of field level extension workers regarding soil analytical results, 'Soil Health Card Program', Use of Upazila Guide for the recommendation of fertilizers on the basis of soil analytical data;
- u) Investigate soil fertility degradation problem, nutrient related problems of crops, soil moisture stress and constraints in crop production etc.

1.2. Organogram of SRDI:

SRDI is headed by Director (Grade 2). Under him there are 4 divisions and 2 research stations.

1.2.1. Development Division

Development division is headed by Chief Scientific Officer and consists of 6 Regional Offices at Divisional level viz. Dhaka, Chittagong, Rajshahi, Khulna, Sylhet and Barisal Regional Office headed by Principal Scientific Officer. There are 5 District Offices headed by Senior Scientific Officer under each Regional Office except Sylhet.

Functions

Planning, co-ordination and supervision of all technical programs and activities of the Regional and District Offices under Development Division. Co-ordinate with allied government and autonomous bodies and other agencies in local and national level program on agricultural development.

1.2.2. Regional Offices

Overall planning, co-ordination and supervision of the technical and administrative affairs of the Regional and District Offices. Co-ordination with the beneficiary agencies like NARS institutes, DAE, BADC, Cotton Development Board, etc. at Regional level planning and execution of agricultural development programs by providing information on soils, land use, land capability, crop suitability, fertility status, etc. Maintaining liaison with Regional Technical Co-ordination Committee and other allied committee/bodies by providing technical assistance, advice, etc. on rational and sustainable use of soil and land resources.

1.2.3. District Offices

Executing and implementing technical activities as per instruction from Head Quarters and Regional Office. Updating Upazila Nirdeshika with basic land and soil information for local level agricultural development planning and extension. Delivering farmers service through crops/cropping pattern based fertilizer recommendation. Technology transfer to farmers through

block demonstration. Assisting beneficiary agencies like DAE, NARS institutes, BADC, Cotton Development Board, etc. with information and advice on matters related to sustainable land and soil resource utilization. Maintaining liaison with District Technical Co-ordination Committee and District Development Co-ordination Committee by providing information on soil and land resources for agricultural and other development planning.

1.2.4. Survey Division

Survey Division is headed by Chief Scientific Officer and consists of 3 Sections viz. Soil and Land Survey Classification sections, Soil Survey Interpretation Section and Soil Correlation Section, each headed by Principal Scientific Officer at Head Quarter.

Functions

Soil Survey planning and co-ordination. Supervision of all technical programs and activities of the component sections. Co-ordination with allied GO/NGO's in national level agricultural development planning. Correlation and classification of soil and land resources at national and international level. Responsible for overall technical activities of the division and field investigation of problems related to soils.

1.2.5. Technical Support Services Division

Technical Support Services Division is headed by Chief Scientific Officer and consists of 5 Sections viz. Central Laboratory, Training Section each headed by Principal Scientific Officer, Cartography Section headed by Senior Cartographer, Publication and Record Section headed by Publication & Liaison Officer and Data Processing & Statistical Section at Head Quarter.

Functions

Planning, co-ordination and supervision of all technical programs and activities of component sections. Review and/or editing all technical reports and maps prepared by using base materials and GIS technology. Co-ordination with allied GO/NGO's in research/investigations involving physical and chemical analyses as well as adaptive research activities relating crops and soils. Storage, analyses and regular updating of soil and land resource database. Maintenance and up scaling of Online Fertilizer Recommendation System (OFRS) and Website management. Responsible for overall technical progress of the division.

1.2.6. Soil Testing Division

Soil Testing Division headed by Chief Scientific Officer and consists of 15 Regional Soil Testing Laboratories and 6 Fertilizer Quality Control Laboratories. Dhaka, Comilla, Rajshahi, Khulna

and Mymensingh Regional Laboratories are headed by Principal Scientific Officer. Ten Soil Testing Laboratories (Barisal, Dinajpur, Bogra, Jamalpur, Noakhali, Kushtia, Faridpur, Jhenaidah, Sylhet and Chittagong) and 6 Fertilizer Quality Control Laboratories are headed by Senior Scientific Officer.

Functions

Analyses of soil, water, plant and fertilizer samples to evaluate problems related to soil fertility, toxicity, salinity and quality control. Based on soil nutrient status and crop requirements farmers are provided with location specific fertilizer recommendation cards (FRC) through static and 12 mobile soil testing laboratories (MSTL). Analytical services are also provided to researcher/beneficiaries.

1.2.7. Research Centers

There are two research centers under direct supervision of Director, SRDI

1. Soil Conservation and Watershed Management Center (SCWMC) is located at Meghla, Bandarban and headed by Principal Scientific Officer. Responsible for generating technology on soil conservation and watershed management in sloping lands of Hilly areas and conducting research on sustainable hill farming.

2. Salinity Management and Research Center (SMRC) is located at Batiaghata, Khulna and headed by Principal Scientific Officer. Responsible for generating data- base on soil and water salinity, identify potential sources of irrigation water, screening of soil tolerant varieties of different crops, innovation of saline soil and water management technologies.

Chapter 2. Activities of Different Sections of Head Quarters

2.1. Soil and Land Classification Survey Section

2.1.0 Soil Survey Planning and Supervision

- Supervision of Photo-interpretative Soil and Landform Map preparation: Kaliganj (Gazipur), Mujibnagar (Meherpur) Upazila
- Supervision of Updating Survey of Kaliganj (Gazipur) and Shirajdikhan (Munshiganj) Upazila



Photo: Supervision of Updating Survey in Kaliganj, Gazipur


2.2.1 Compilation of district Soil and Landform Map

Rationale:

- Popularize the use and soil and landform data among Stakeholders
- Enhance district level agricultural development planning with land and soil data
- Support in planning district agricultural rehabilitation program

Methodology

- Upazila Soil and Landform Maps (1:50,000) were digitized and scale was converted to 1:125,000.
- Edge matching of each mapping unit was done.
- A common legend for the district map was constructed accommodating Upazila map legends.



Map 1: Compiled Soil and Landform Map, Dhaka District

Digital Soil Map Preparation for Rangpur District Applying World Reference Base (WRB)

Objectives: The main objectives of digital soil map preparation for Rangpur district applying World Reference Base (WRB) are to:

- ❑ Correlate soils of Rangpur district with WRB so as to harmonize the respective soils with globally recognized soil classification and naming system.
- ❑ Simplify resource management domain for planning the best possible way to transfer agro-technology relevant with the local soil environment.
- ❑ Create inventory of soil resources which could be the tool to identify climate smart agriculture adaptive to particular soils.

Methodology




Figure 1. Sketch representation of the method for digital mapping applying WRB principles

Basic Principles of WRB

- ❑ Step – 1: Detecting diagnostic horizon, properties and materials
- ❑ Step – 2: Allocating the soil to a Reference Soil Group (RSG), e.g. Acrisols, Gleysols etc.
- ❑ Step – 3: Allocating the qualifiers, such as Eutric, dystric etc.

Construction of Soil Map Legend (in WRB)

Map legend describes the map units. A map unit consists of

- ❑ A dominant soil (covering $\geq 50\%$) of the soil cover only or
- ❑ A dominant soil plus a co-dominant soil (covering $\geq 25\%$ and $\leq 50\%$) of the soil cover and/or one or more associated soils (covering $\geq 5\%$ and $\leq 25\%$) of the soil cover or
- ❑ Two or three co-dominant soils or
- ❑ Two or three co-dominant soils plus one or more associated soils

Findings


Total number of mapping units (in 7 upazilas) at Upazila scale: **71**

Total number of mapping units after harmonization of mapping unit considering physiography, land type and soil at district scale: **19**

Total number of mapping units after harmonization applying WRB principles of map legend at district scale: **9**

Soil classification applying WRB

Soil Classes in WRB	Soil Group /Series	Total Area (%)
Haplaic Acrisols	Kashimpur, Belabo	6.7
Eutric Cambisols	Domar, Pirgachha, Palashbari	15.8
Eutric Gleysols	Noadda, Chandra, Nijhuri, Sahapur, Ekdala, Lauta, Khilgaon, Jamun, Gangachara, Kaunia, Bhimpur, Indrail, Lashkara, Kakina, Kachna, Chilmari	71.65
Eutric Fluvisols	Manda, Shaghatta, Tista Sandy Alluvium, Tista Silty Alluvium	5.85



Map 2: Soils of Rangpur District, WRB

Survey and report preparation regarding siltation on agricultural fields in flood-affected areas

The survey conducted and report prepared on sand/silt deposition on flood-victim areas of agricultural fields jointly by SRDI and BARC after field survey during November 2017.

Effects of Brick kiln on Environmental Pollution and Degradation of Agricultural Land

Summary findings

A total of 121 brick kilns have been surveyed through this project. The same number of the owners of the associated brick kilns has been interviewed. The survey area covered 86 Upazilas (sub-districts) under 48 districts throughout Bangladesh. Most of the brick kilns are using coal as fuel. Forest woods were used in 18% of the brick kilns. Surveyed brick kilns are situated within 3-20 km from the Upazila Headquarters and 2m-2km from the nearest homesteads. The average distance between brick kilns and nearest homesteads is about ½ km (570 m) and the distance between brick kilns and Upazila HQs is about 7.36km. About 58% (69 Nos.) brick kilns were approved by the competent authority; 18% (21 Nos.) had no approval. This information was not disclosed by 21% brick kiln owners. The distance and locations maintained by brick kiln owners show clear violation of Brick Making and Establishment of Kilns (Control) Law 2013. Nearby farmers reported adverse effect of brick kilns on homestead plantation and field crops. They also reported that bearing of fruit trees is hampered; immature fruits are damaged, less fruiting in vegetables.

The total production of 121 brick kilns per annum is 31.70 million bricks. The capacity of the surveyed brick kilns ranges from 0.45 million to as high as 0.78 million bricks per year. The average capacity was 2.89 million bricks per year. Bangladesh produces around 50 billion bricks per year. Use of quality topsoil of agricultural land as raw material for bricks is a substantial environmental and economic concern, which has led the government to prohibit previously profitable brick exports to India as per law of the land.

Mostly fertile topsoil of agricultural land is exploited for making bricks. On an average 3 kg soil is required to make a brick. Thus, 150 billion kg soil is required to produce 50 billion bricks per annum in Bangladesh. One hectare land up to furrow slice i.e. 0-10 cm contains 2×10^6 kg of soil. Bangladesh cannot afford losing huge amount fertile topsoil every year.

2.2 Soil Survey Interpretation Section

Responsibilities

- 1) Soil Survey Interpretation Section is responsible for planning, supervision and execution of soil survey interpretation activities for various beneficiaries engaged in agricultural development
- 2) Generating basic data on soils, land capability and crop suitability for preparation of short and long term agricultural development plans/projects.
- 3) Interpretation of soil database for location specific crop suitability assessment and processing of soil survey data for developing and updating GIS based data bank.

Achievements (2017-18) are shown under following heads

Updating Soil Survey program

Upazila Land & Soil Utilization Guide Preparation

Sustainable Land Management

Documentation of SLM technologies through WOCAT software

Annual Performance agreement

NIS (National Integrity strategy) related Activities

Sustainable Development Strategy

Technical Specification Activities

Innovation Activities

Other Activities

Conducting Field Survey for Updating Upazila Nirdeshika

1. KaligongUpazila (21,316 hectare) under Gazipur district

Physiography: 1) MadhupurTract, 2) Old Brahmaputra floodplain

(3) Meghna estuarine floodplain

2. SirajdikhanUpazila under Munshigong district.

Change of land type over the years: Kaliganj Upazila

Previous (2000)			Present (2018)			Change
Land type	Area(ha)	(%)	Land type	Area	(%)	Increase/Decrease
HL	6,679	29.0	HL	5,877	27.6	-(1.4)
MHL	5,682	26.5	MHL	5,548	26.0	-(0.5)
MLL	1,970	9.3	MLL	2,235	10.5	+(1.2)
LL	2,883	13.5	LL	2,710	12.7	-(0.8)
VLL	2,094	9.8	VLL	2,053	9.6	-(0.2)
Misc.	2,535	11.9	Misc.	2,893	13.6	+(1.7)
Total	21,316	100	Total	21,316	100	-

Present cropping pattern in Kaliganj Upazila (2018)

SL	Cropping patterns	Area (ha)	Percentage
1.	Perennial crops	1,897	8.9
2.	Annual crops (Pine apple/ Ginger/Turmeric	916	4.3
3.	Rabi vegetables- Kharif vegetables	532	2.5
4.	Rabi crops- Boro-T. Aman	2877	7.2
5.	Mustard-Boro- T.Aman	1683	7.9
6.	Rabi crops- jute- T. Aman	788	3.7
7.	Boro-Fallow-Fallow	3602	16.1
8.	Boro-Fallow -T. Aman	2685	12.6
9.	Mustard-Boro-Fallow	1236	5.8
10.	Others cropping patterns		
11.	Miscellaneous (Homestead, River, Ponds and Others)	2893	

Previous cropping pattern in Kaliganj Upazila (2000)

SL	Cropping patterns	Area (ha)	Percentage
1.	Perennial crops	2,757	12.9
2.	Annual crops (Pine apple/ Ginger/Turmeric	793	3.7
3.	Rabi vegetables- Kharif vegetables	375	1.7
4.	Fallow- Kharif vegetables	1554	7.3
5.	Rabi crops/rabi vegetables- B. Aus/ jute- Fallow	1542	7.2
6.	Mustard-Boro- T.Aman	381	1.7
7.	Rabi crops- B. Aus/jute- T. Aman	1266	5.9
8.	Fallow- B. Aus- T. Aman	1288	6.0
9.	Boro-Fallow-Fallow	1545	7.2
10.	Boro-Fallow -T. Aman	1545	7.2
11.	Mustard-Boro-Fallow	670	3.1
12.	Miscellaneous(Homestead, River, Ponds and Others)	2535	

Changes in land use: Fallow- Kharif Vegetables; Rabi Crops/Rabi Vegetables- B. Aus/ jute-Fallow; RC- B. Aus/jute- T. Aman patterns have become marginalized. On the other hand, the growing area of Mustard, Boro as well as T. Aman has increased significantly.

Upazila Nirdeshika Report Preparation

Updated Upazila Nirdeshika of Dhamrai Upazila under Dhaka has been submitted to Upazila Nirdeshika Cell, HQ, Dhaka.

APA activities

Drafting and finalization of APA 2018-19 as per format supplied by MoA/CD, Govt. of Bangladesh

Supervision of Drafting and finalization of APA 2018-19 for field offices Three Quarterly and One Half yearly Monitoring

Progress report of APA 2017-18 submitted to MoA.

Document collection, preservation and submission to MoA as well as Expert pool of MoA as evidence of activities performed according to APA2017-18 during half yearly performance evaluation by Expert pool of MoA.

1st, 2nd, 3rd & final draft of APA for 2018-19 submitted to MoA.

One day training program is being arranged for the field officers on Drafting APA 2018 -19 .

Attended the meeting & Lab training at MoA/CD on regular basis as focal point of SRDI APA.

National Integrity Strategy (NIS):

Drafting and finalization of NIS work plan 2018-19 as per format supplied by MoA/CD.

Supervision of Committee formation, Drafting and finalization of NIS work plan 2018-19 as per format supplied by MoA/CD for field offices.

Three Quarterly Monitoring Progress report of NIS 2018-19 submitted to MoA.

One day training programmed is imparted for the field officers on NIS.

Attended the meeting & training at MoA regularly as focal point of SRDI NIS Committee.

Other activities:

Performing secretarial job for Innovation team of SRDI.

Participating Innovation training conducted by a2i, PMO, MoA.

Data compilation: Meteorological data compilation from 1985 to 2016

Monthly progress report preparation

Training Received: 1. Training on climate smart Agriculture

2. Training on documentation SLM technologies through WOCAT software
3. Training on Integrated Budgeting & Auditing System
4. Training on service simplification
5. Training on Innovation


2.3 Soil Correlation Section

2.3.1 Function of Soil Correlation Section

1. Planning, supervision & execution of soil series & other taxonomic units
2. Maintenance of uniform standard of methodology on soil survey works and & records of soil information
3. Correlation of soil surveys done by other agencies / consultancy
4. Development & maintenance of the soil museum

2.3.2 Achievements

The information given on Monolith boards has been redesigned and proposed as follows.



Exhibits in the Soil Museum

Physiography-wise soil monoliths and soil correlation boxes collected, preserved, re-arranged, and displayed in the museum are:

- Physiography --- 11 (out of 15)
- Soil Monoliths --- 50 (48 SRDI & 2 BARC) out of 476 soil series (453 soil series, 23 different river alluviums)
- Soil Monoliths, recently collected - 6
- Correlation Box with soil series display-1178
- Correlation box (blank) - 1050 (50 new)




Figure 2: Graphical representation of the Soil Monoliths Exhibits in the soil museum

2.4 Training Section

Training Section arranged various types of local training for human resource development for the officers and staffs of different levels as part of its activities for the year 2017-2018. This includes in-service training of the freshly recruited officers on various aspects of technical activities of the institute, training on administrative, office management, ICT and financial management for officers and staffs. This section made arrangement for foundation training of newly recruited officers also made arrangement for the officers and staffs of SRDI to take part in different training programs of local and international level as well as in different workshops and seminars for a number of occasions held in different organizations upon their invitations. Training section mainly provided logistic and technical support for smooth functioning of those training programs. A good number of officers participated at Radio talk in different agricultural programs

of Bangladesh Betar. Training Section communicated with Bangladesh Betar and proposed names of officers with titles of radio talk to be broadcasted in collaboration with AIS and DAE.

In-house Training Plan & Achievement-2018

Grade	No. of Officers & Staff	Target		Achieved (up to May, 18)	% of Achievement against target to May'18
		Per person	Total		
2-9	154	60	8,640	3,280	95
10-12	18	60	1,080	365	96
13-17	217	60	13,020	4,375	89
18-20	141	60	4,460	3,075	93

Activities done by the Training Section for the year 2017-2018

(i) Higher Education

Sl. No.	Degree	On-going	No. of Officers				Remarks
			Obtained Scholarship	Received permission	Requested for permission	completed	
1	PhD	12	13	3	10	4	-

(ii) Foundation Training for 6 Months

Sl. No.	No. of Trainees		
	BCS (Ag.) Cadre	Non-Cadre	Total
1	6	-	6

(iii) Special Training Arranged by Training Section

Sl. No.	Courses	No.	Duration
1	Training on Arc GIS Software for Better GIS Activities in Producing Maps and Reports	20	6 days
2	Training on Preparation of Upazila Nirdeshika with Soil and Land form Map Using Updated Aerial Photographs	33	5 days
3	Training on Skill Development through Laboratory Analytical Methodology	25	2 days
4	Training on National Integrity and Strategies	30	2 day
5	Training on National Integrity and Strategies & Annual Performance Agreement	25	2 days

(iv) In-house Training Arranged by Training Section (for Staffs of Head Office)

Formal Monthly in-house training

Grade	No. of Officers/Staffs (every month)	Duration
10-12	14	1 day
13-17	63	
18-20	43	

In-house Training Topics

- The Public Employee's Discipline (Punctual Attendance), 1982
- The Govt. Servants (conduct) Rules, 1979
- The Govt. Servants (Discipline and Appeal)
- The General Provident Fund Rules, 1979.
- Keeping Service Records
- Store Management
- Public transport purchase, maintenance and repairing
- Use of office equipment and other electronic items
- Cleanliness and dress-up
- Office Security
- Drafting letters and notes with special emphasis to Promito Bangla Bananer Nyom)
- National Integrity Strategy
- Administrative Tribunal Act, 1980 & Rule, 1982
- Digital filing
- Foreign Tour, Lien
- Fundamental right as per constitution and service rule
- Liveries, allocation and conditions
- Food Based Nutrition
- APA, Citizen Charter
- Maintenance and repairing of vehicles, driving rules and regulations
- Delegation of Financial Power
- Advances
- General Principles of Seniority
- ACR
- Treatment facilities for govt. Servants

(v) Training/Seminar attended abroad

Sl. No.	Courses	No.	Duration	Country
1	SOIL ATLAS OF ASIA, Editorial Board Kick off Meeting	1	3 days	Philippines
2	Lab Management for SEALNET	1	5 days	Indonesia
3	Sustainable Diversification in Tropical Crop-Livestock Systems in Asia: Biological Nitrogen Fixation by Forage Legumes	2	5 days	China
4	Training Workshop on Eco-friendly Agricultural	1	25 days	Korea

	Technology			
5	International Symposium on Soil Health and Sustainable Development and Fourth Asian Soil Partnership Workshop	1	4 days	China
6	Seminar on China's Experience in Agricultural Developing Countries	1	21 days	China

Training/Seminar nominated for abroad

Sl. No.	Courses	No.	Duration	Country
1	Resource Recovery Option of Faecal Sludge Management (FSM) Value Chain	1	4 days	Thailand
2	Training Program on Integrated Nutrient Management for Improving Soil Health and Crop Productivity	1	10 days	India
3	International Conference on Earth Science and Geo-science 2018	2	2 days	Netherlands

(v) Local Training Programs attended by officers

Sl.	Courses	No.	Duration	Institute
1	Training on Forestry and Forestry Technology for Professionals	2	3 days	BARC
2	Climate Change, Carbon Sequestration and Adaptation Strategies	3	3 days	
3	Intellectual Property Rights (IPR) and Technology Management	3	3 days	
4	Project Development and Management	1	5 days	
5	World Overview of Conservation Approaches and Technologies	2	2 days	
6	Commemoration of the 33 rd SAARC Chartered Day and Celebration of International Year of Pulse	5	1 day	
7	Capacity Development for Advanced Agricultural Research Methodology and Scientific Report Writing	2	7 days	
8	Technical Report Writing and Editing	2	5 days	
9	Financial Management Course	5	5 days	NATA
10	Modern Office Management	2	5 days	
11	Change Management through ICT	1	4 days	
12	Food Security	4	5 days	
13	Disaster Management	1	5 days	
14	Public Procurement Procedure	6	10 days	
15	Climate Smart Agriculture	6	5 days	
16	Project Appraisal and Formulation of Project Proforma DPP	1	5 days	
17	Training of Trainers on Teaching Methods and Techniques	4	5 days	
18	Innovation in Public Service	2	5 days	
19	Advanced ICT Management	4	10 days	

Sl.	Courses	No.	Duration	Institute
20	Implementation of SDG, NIS and APA	2	2 days	
21	Good Governance	1	5 days	
22	Commercial Farm Management	1	5 days	
23	Creating Congenial Environment in Innovation for the Members of Innovation	1	6 days	MoA
24	Implementation of Fifth Five Year Plan and SDG	1	5 days	
25	Project Development and Planning Process Management	1	4 days	MoF
26	Training on e-GP	4	2 days	
27	Training on Budget Preparation-1	3	4 days	
28	Budget Management Specialist	4	3 days	
29	iBAS++ Training in Accounting and Budgeting	2	1 day 5 times	BMD
30	Use of Satellite Data Products for Drought Monitoring	1	5 days	
31	The Introduction to Electromagnetic Induction Techniques for Soil Salinity Investigations	1	3 days	NUMAN
32	Orientation and Sub-project Selection	2	1 day	LGED
33	Statistical Analysis R-software	2	5 days	BARI
34	Financial Management	1	10 days	BIAM
35	Climate Change Issues and its Adaptation	2	5 days	BARD
36	Administrative and Financial Management	1	15 days	
37	Project Planning, Implementation, Monitoring and Evaluation	1	10 days	
38	Development Management	1	5 days	
39	Research Methodology	3	15 days	GTI, BAU
40	Professional GIS Training Course	2	5 days	DU
41	TIBAS Training in Accounting and Budgeting	5	10 days	FIMA
42	Training of Trainer on Food Based Nutrition	8	5 days	BIRTAN

(vi) Local Training Programs attended by staffs

Sl.	Courses	No.	Duration	Institute
1	Public Procurement	3	2 days	RPATC
2	Discipline, Ethics and Etiquette	3	12 days	
3	Basic Office Management	7	1 day	
4	ICT and e-Governance Management Course	2	15 days	
5	Computer Literacy and English Language Course	2	3 days	
6	Fundamental Financial Rules	4	1 day	
7	Computer Application and English Language	2	12 days	
8	Office Automation for Organizational Development	1	15 days	
9	Staff Development	2	3 days	
10	Communicative English	2	12 days	

(vii) Local Workshop/Seminar attended

Sl.	Title	No.	Duration	Institute
1	Advanced Training Workshop on Piloting of the Poverty Environment Accounts (PEA)	3	3 days	BBS
2	Exhibition and Regional Seminar on Agricultural Mechanization to Commemorate 33 rd SAARC Charter Day	2	1 day	BARC
3	Workshop on APASS	2	1 day	
4	Strategic Management Plan for NATA	2	1 day	NATA
5	Training Need Assessment	4	1 day	
6	Use of Research Archive Software	30	1 day	SRDI
7				
8	Computer Application and English Language	2	12 days	
9	Office Automation for Organizational Development	1	15 days	
10	Staff Development	2	3 days	
	Communicative English	2	12 days	

Training Need Assessment

Training Need					
Management			Technical		
Training Title	Grade	Duration	Training Title	Grade	Duration
ICT Management	9-4	3 weeks	Natural Resource Management	9-4	4 weeks
Modern Office Management	9-5	2 weeks	Sustainable Land Management and Poverty Reduction	9-5	2 weeks
Financial Management	9-4	2 weeks	Studies on Soil and Plant relationship	9-5	2 weeks
Procurement, PPR	9-4	3 weeks	Integrated Nutrient Management	9-3	3 weeks
Database Management	9-3	4 weeks	Remote Sensing and GIS	4-2	3 weeks
Policy and Strategies for Sustainable Soil and Land Management	9-3	1 weeks	Soil Carbon Sequestration	9-5	2 weeks
Training on GIS Software	9-3	3 weeks	Use of GIS in Soil and Land Management	9-5	2 weeks
Institutional Capacity Building	4-2	2 weeks	Soil Fertility and Productivity Management	9-5	3 weeks

Other Activities Done by Training Section

- Preparing DPP and all corresponding works of the projects submitted to MoA
- Successful implementation of E-Filing Program
- APA & SDG Action Plan Activities
- Preparing different Reports, Booklets, Directory etc.
- Innovation activities of SRDI
- Execution of PM's commitment
- Procurement work of SRDI
- Organogram, recruitment rules, different cases etc.

Training Plan

- Making arrangement of all types (mentioned earlier) of local and foreign trainings as per instruction of authority
- Strengthening training activities
- Arrangement of in-service training for the newly recruited officers
- Arranging short training course on Establishment Manual-2008, Jatio Shuddhachar, ICT, Nutrition, Office Management etc.
- Arranging some refreshers course
- Arrangement of radio talks on different topics related to soil fertility, soil management etc. for Bangladesh Betar, BTV and other Channels.

2.5 Data Processing and Statistical Section

Activities of DPS Section

- Engaged in planning, organizing and execution of GIS related works
- Digitizing, preparation and printing of different types of thematic maps.
- DPSS is responsible for storage, maintenance and security of database on soil and land resources and other information

Major Type of Works

1. GIS related
2. ICT related
3. Others

GIS Related Works

- Preparation of geo-referenced and geo-projected database
- Map Preparation & Printing-
 1. AEZ and Soil Salinity Map with Fertilizer Factory & Buffer Godown Location

2. Flood Prone Area Map
3. Soil Organic Carbon Status Map
4. AEZ Map in Bengali
5. AEZ- Old Himalayan Piedmont Plain Map
6. Northern & Eastern Hills Map
7. AEZ- Old Brahmaputra Floodplain Map
8. AEZ- Ganges Tidal Floodplain Map
9. Madhupur Tract Map
10. Agro Ecological Regions of Dhaka Division Map
11. Agro Ecological Regions of Dhaka District Map
12. Agro Ecological Regions of Barisal Division Map
13. AEZ Map of Chandpur District
14. Salinity Map of Amtali Upazila (Barguna District)
15. Salinity Map of Monpura Upazila
16. Salinity Map of Kalapara Upazila (Patuakhali District)
17. Barind Tract and Modhupur Tract of Bangladesh
18. Organic Matter Status Map of Mymensingh Sadar Upazila




Fig.3 AEZ and Soil Salinity Map with Fertilizer Factory & Buffer Godown Location




Fig.4 Flood Prone Area Map based on land type data of Upazila Nirdeshika




Fig.9 AEZ- Old Brahmaputra Floodplain Map




Fig.10 AEZ- Ganges Tidal Floodplain Map




Fig.11 Madhupur Tract Map




Fig. 12 AEZ Map of Dhaka Division




Fig. 13 AEZ Map of Dhaka District




Fig. 14 AEZ Map of Barisal Division




Fig. 15 AEZ Map of Chandpur District




Fig.16 Salinity Map of Amtali Upazila (Barguna District)




Fig.17 Salinity Map of Monpura Upazila




Fig.18 Salinity Map of Kalapara Upazila (Patuakhali District)




Fig.19 Barind Tract and Modhupur Tract of Bangladesh




Fig. 20 Northern & Eastern Hills of Bangladesh




Fig.21 Maps of Organic Status in different time (1990 and 2015) showing the increasing trend of Organic Matter in soil of Mymensingh Sadar Upazila

ICT Related Works

1. Server & LAN management

- Proper monitoring, maintenance and trouble shooting of the server and internet related devices (server, router, bandwidth controller, switch etc.) of SRDI
- At present there are 86 internet connection in SRDI head office

2. Data Processing & Uploading

- Soil Chemical data are generalized, processed and prepared for uploading.
- This uploaded data is used in the software for Online Fertilizer Recommendation System (OFRS).

3. Others Support

- Hardware maintenance and trouble shooting of different computer of all section of SRDI.
- Take part with proper logistic support of different agricultural fair.
- Technical assistance and necessary support provided to develop Soil Organic Carbon Map of Bangladesh with the help of UNCCD.
- Technical assistance and necessary support provided to prepare different on demand report from MoA, BARC and others.

- Technical assistance and necessary support provided for sharing GIS meta data for Bangladesh Geographical Information Platform (BGIP) organized by BBS.
- Upazila survey of Kaliakair and Rupganj Upazila are carried out by two officers of DPS Section.

Plans

- We are trying to retrieve our old database (where projection file is missing) for working with concurrent GIS software. For this we need some expert technical assistance.
- We are updating our soil nutrients shape file by digitizing point file with necessary attributes for updating soil nutrient maps.
- Different types of map will be prepared as per demand.

2.6 Cartography Section

1. Map Digitizing

Name of Map	Division/District/Upazila Name	Map Scale	Nos
Cotton Suitability Area Map	Poba, Islampur, Muksudpur, Potnitola, Barisal Sadar, Kalaroya, Nalitabari, Comilla Sadar Dakshin, Nageshwari Upazila	1:50000	9
Soil and Landform Map	Muksudpur, Tongibari, Patharghata Upazila	1:50000	3
Salinity Map	Khulna District	1:25000	1
Sketch Map	Batiaghata Office Area		1
Mouza wise Upazila Map	Tongibari, Manikganj, Narsingdi Upazila	1:50000	3

2. Map Tracing

Name of Map	Division/District/Upazila Name	Map Scale	Nos
Soil and Landform Map	Baralekha, Kumarkhali, Nalitabari, Islampur, Bhuyapur, Dhamrai, Nachole, Pagla, Bahubal, Bancharampur, Chhatak Upazila	1:50000	11
Union wise Soil and Landform Map	Belabo (4), Kapasia (3), Dhamrai (1), Singair (2), Tongibari (12)	1:50000	22
Mouza wise Upazila Map	Various Upazila Map	1:4000	18

3. Map Checking & Correction

Name of Map	Upazila/Union Name	Map Scale	Nos
Various Map	Various Upazila and Union Map	1:50000	45

4. Map Printing

Name of Map	Upazila/Union Name	Map Scale	Nos
Soil and Landform Map	Various Upazila and Union Map	1:50000	240

4. Map Colouring

Name of Map	Upazila/Union Name	Map Scale	Nos
Upazila and Union	Soil and Landform Map of various Upazila and Union	1:50000	125

5. Area Calculation

Name of Map	Upazila/Union Name	Map Scale	Nos
Soil and Landform Map	Various Upazila and Union Map	1:50000	20

6. Collection of Base Materials from Survey of Bangladesh

Item	Upazila Name	Map Scale	Nos
Arial photo	36 Upazila	1:25000	875
Arial photo	27 Upazila's latest Arial photo collection from SoB under processing	1:25000	410 (approx.)
Topo Map	-	1:25000	160
Topo Map	Latest Topo Map collection from SoB under processing	1:25000	700 (approx.)

2.7 Publication and Record Section

Publication and Record Section of the Technical Support Services Division is responsible for printing, publication and distribution of soil survey and other technical reports and their overall maintenance, to keep liaison with outside agencies for the above mentioned purpose and to assist the authority in technical & administrative support on different aspect.

1. Sale of Upazila Nirdeshika

Name of Organization	Type	Purpose	Quantity
BWDB	Autonomous	Project	39
Dhaka University	Autonomous	Academic	5
Jahangir Nagar University	Autonomous	Academic	02
Agrarian Research Foundation	NGO	Research	19
Blue Gold Program	NGO	Research	07
Pro Soil Foundation	NGO	Research	48
Personal	Student	Academic	37
Total			157

2. Other Activities

- Collection of 50 copies of Books and Journals
- Arrangement of Nomination for
 - Bangladesh National Agriculture Award
 - Ekushe Podok
 - Independent Award
 - Begum Rokeya Award
- Distribution of Posters-30 and Booklets on Identification of Adulterated Fertilizers-20
- Readers service- 300 readers

2.8 Central Laboratory

1.2 Introduction

Central Laboratory is operated under Support Service Division in the headquarter of SRDI and the functions are- execution of physical and chemical analyses of soil samples collected by the survey and development divisions and also samples sent by beneficiary organizations including DAE, NARS organizations, universities, farmers, entrepreneurs and NGO's. Central Laboratory also conducts chemical analysis of water, plant and fertilizer samples. It is also responsible for planning and supervision of basic and adaptive research works such as soil degradation, crop and soil moisture relationship studies, variation of nutrient status of different soils, soil toxicity, etc. Mobile soil testing laboratory named Jamuna, is also operated by this Laboratory for the awareness of farmers regarding judicious application of fertilizers and inspire the farmers using soil test base balanced fertilizer application. Central laboratory has generated technologies like identification of adulterated fertilizers at field level without laboratory test. Quality control activities on different fertilizers are one of the most important works of SRDI which is mostly done by Central Laboratory.

1.2 Major activities of Central Laboratory

1. To determine soil fertility and fertility status for the recommendation of soil test base balanced fertilizer doses for different crops;
2. To analyze soil samples for preparing Upazilla Land and Soil Resources Utilization Guide and Union Sahayika;
3. To analyze water and plant samples received from different stakeholders;
4. To provide farmers soil sample analysis through Mobile Soil Testing Laboratory and
5. To determine nutrient content of fertilizers under the quality control activities of the government.

Beside the above activities Central Laboratory provides advisory service to the farmers, impart training programs and conducts research programs on soil and fertilizer management.

1.3 Soil analyses

Central Laboratory analyzed a total of 2166 soil samples under different programs namely Farmer's Service through Static Laboratories and Mobile Soil Testing Laboratories, Preparation of Land and Soil Resources Utilization Guides, salinity survey and supporting research activities of different institutes.

1.3.1 Materials and Methods

A total of 137 soil samples were received from the farmers of 35 districts under 8 divisions. To determine the nutrient content and evaluate the fertility status of farmers samples both micro and macro nutrient elements as well as pH and organic matter content of the soil were determined. Worldwide established standard methods were followed to analyze different nutrient elements. Nitrogen was determined by Micro Kjeldahl method, phosphorus determined by Bray and Kurtz method (if pH is <7.0) or Olsen method (if pH is >7.0) for, sulfur determined by Turbidimetric method and organic matter determined by Walkley and Black Wet oxidation method etc.

1.3.2 Results and discussion

The results and discussions are made below division wise.

Table 1: Soil samples received from farmers of different divisions and analyzed at Central Laboratory in 2017-2018

Serial No.	Division	Number of Samples
1.	Dhaka	44
2.	Chattogram	31
3.	Rajshahi	11
4.	Khulna	13
5.	Barishal	12
6.	Rangpur	14
7.	Sylhet	2
8.	Mymensingh	10
	Total	137

Note: The results found here should not be taken as the country's profile. However, it is indicative of the overall fertility status of the plots of the farmers, clustering in different divisions, who sent their samples to Central Lab for analysis.

Soil pH

It was observed that pH of the soil samples of Dhaka division ranges from 4.0 - 7.8 of which 3% samples are very strongly acidic, 23% samples are strongly acidic, 41% samples are slightly acidic, 20% samples are neutral and 14% samples are slightly alkaline. In Chattogram division soil pH ranges from 4.0 - 8.6 where 2% samples are very strongly acidic, 45% samples are strongly acidic, 16% samples are slightly acidic, 10% samples are neutral, 16% samples are slightly alkaline and 10% samples are strongly alkaline. In Rajshahi division soil pH ranges from 5.4 - 6.9 of which 9% samples are strongly acidic, 55% samples are slightly acidic and 36% samples are neutral. In Khulna division soil pH ranges from 6.4 - 7.9 of which 8% samples are slightly acidic, 46% samples are neutral and 46% samples are slightly alkaline. In Barishal division soil pH ranges from 5.2 - 7.7 of which 8% samples are strongly acidic, 42% samples are slightly acidic, 42% samples are neutral and 8% samples are slightly alkaline in fertility status. In Rangpur division soil pH ranges from 4.8 - 6.5 of which 36% samples are strongly acidic and 64% sample are slightly acidic. In Sylhet division soil pH of farmers field ranges from 4.3 - 4.4 of which 100% samples are very strongly acidic and in Mymensingh division soil pH ranges from 4.6 - 6.9 of which 40% samples are strongly acidic, 40% samples are slightly acidic and 20% samples are neutral in fertility status.

Organic matter

The organic matter content in the soil samples of Dhaka division ranges from 0.20 % – 4.24 % where 27% samples are very low, 32% samples are low, 30% samples are medium, 7% samples are high and 4% samples are very high in fertility status. Organic matter in soil samples of Chattogram division ranges from 0.20 % – 2.69% of which 48% samples are very low, 35% samples are low and 17% samples are medium in fertility status. Soil samples from Rajshahi division organic matter content ranges from 0.54 % – 2.55% of which 18% samples are very low, 45% samples are low and 37% samples are medium in fertility status. The organic matter content of soil samples of Khulna division ranges from 0.54% – 2.42% of which 23% samples are very low, 54% samples are low and 23% samples are medium in fertility status. The organic matter content of soil samples of Barishal division ranges from 0.87 % – 4.17% of which 8% samples are very low, 33% samples are low, 50% samples are medium and 9% samples are high in fertility status. The organic matter content of soil samples of Rangpur division ranges from 1.55 % – 1.82% of which 50% samples are low, 43% samples are medium and 7% samples are

high in fertility status. The organic matter content of soil samples of Sylhet division ranges from 1.14 % – 3.77% of which 50% samples are low and 50% samples are medium in fertility status. The organic matter content of soil samples of Mymensingh division ranges from 0.67 % – 2.89 % of which 10% samples are very low, 30% samples are low, and 60% samples are medium in fertility status.

Total nitrogen

It is observed that the total nitrogen content of soil samples of Dhaka division ranges from 0.010 % – 0.414% of which 57% samples are very low, 32% samples are low, 7% samples are medium, 2% samples are optimum and 2% samples are high in fertility status. The total nitrogen content of soil samples of Chattogram division ranges from 0.010 % – 0.135 % of which 84% samples are very low and 16% samples are low in fertility status. The total nitrogen content of soil samples of Rajshahi division ranges from 0.027 % – 0.128 % of which 64% samples are very low and 36 % samples are low in fertility status. The total nitrogen content of soil samples of Khulna division ranges from 0.027 % – 0.121 % of which 77% samples are very low and 23% samples are low in fertility status. The total nitrogen content of soil samples of Barishal division ranges from 0.044 % – 0.209 % of which 42 % samples are very low, 50% samples are low and 8% samples are medium and the total nitrogen of soil samples of Rangpur division ranges from 0.057 % – 0.189 % of which 50% samples are very low, 43% samples are low and 7% samples are medium in fertility status. The total nitrogen content of soil samples of Sylhet division ranges from 0.078 % – 0.091 % of which 100% samples are very low and total nitrogen content of 10 soil samples of Mymensingh division ranges from 0.034 % – 0.330 % of which 40% samples are very low, 50% samples are low and 10% samples are optimum in fertility status.

Phosphorus

The phosphorus content of soil samples of Dhaka division ranges from 0.43 ppm – 176.77 ppm of which 55% samples are very low, 9% samples are low, 11% samples are medium, 5% samples are optimum, 2% samples are high and 18% samples are very high in fertility status. The phosphorus content of soil samples of Chattogram division ranges from 0.41 ppm – 36.46 ppm of which 87% samples are very low, 7 % samples are low and 3% samples are medium in fertility status. The phosphorus content of soil samples of Rajshahi division ranges from 2.18

ppm – 97.14 ppm of which 55% samples are very low, 9 % samples are medium and 9% samples are optimum in fertility status. The phosphorus content of soil samples of Khulna division ranges from 0.13 ppm – 90.65 ppm of which 54% samples are very low, 8% samples are low, 15% samples are medium and 23% samples are very high in fertility status. The phosphorus content of soil samples of Barishal division ranges from 0.53 ppm – 113.59 ppm of which 50 % samples are very low, 8% samples are low, 8% samples are high and 34% samples are very high in fertility status. The phosphorus content of soil samples of Rangpur division ranges from 4.21 ppm – 92.55 ppm of which 7% samples are very low, 36% samples are medium, 14% samples are optimum, 7% samples are high and 36% samples are very high in fertility status. The phosphorus content of soil samples of Sylhet division ranges from 1.43 ppm – 7.56 ppm of which 50% samples are very low and 50% samples are low and the phosphorus content of Mymensingh division ranges from 1.15 ppm – 33.02 ppm of which 60% samples are very low, 20% samples are low and 10% samples are medium in fertility status.

Potassium

The potassium content of soil samples of Dhaka division ranges from 0.10 meq /100gm soil – 1.54 meq /100gm soil of which 45% samples are low, 18% samples are medium, 14% samples are optimum and 23% samples are very high in fertility status. The potassium content of soil samples of Chattogram division ranges from 0.06 meq /100gm soil – 0.35 meq /100gm soil of which 29% samples are very low, 29 % samples are low, 26% samples are medium and 16% samples are optimum in fertility status. The potassium content of soil samples of Rajshahi division ranges from 0.11 meq /100gm soil – 0.60 meq /100gm soil of which 27 % samples are low, 27% samples are medium, 27% samples are optimum, 10% samples are high and 9% samples are very high in fertility status. The potassium content of soil samples of Khulna division ranges from 0.20 meq /100gm soil – 0.42 meq /100gm soil of which 38% samples are medium, 54 % samples are optimum and 8% samples are high in fertility status. The potassium content of soil samples of Barishal division ranges from 0.15 meq /100gm soil – 0.40 meq /100gm soil of which 25% samples are low, 42% samples are medium, 17% samples are optimum and 16% samples are high in fertility status. The potassium content of soil samples of Rangpur division ranges from 0.09 meq /100gm soil – 0.40 meq /100gm soil of which 22% samples are very low, 57% samples are low, 14% samples are medium and 7% samples are high in fertility status. The potassium content of soil samples of Sylhet division ranges from 0.08 meq

/100gm soil – 0.09 meq /100gm soil of which 100% samples are very low and the potassium content of 10 soil samples of Mymensingh division ranges from 0.07 meq /100gm soil – 0.34 meq /100gm soil of which 30% samples are very low, 60% samples are low and 10% samples are optimum in fertility status.

Sulfur

The sulfur content of soil samples of Dhaka division ranges from 1.23 ppm– 223.06 ppm of which 41% samples are very low, 20% samples are low, 9% samples are medium, 7% samples are optimum, 9% samples are high and 14% samples are very high in fertility status. The sulfur content of soil samples of Chattogram division ranges from 0.91 ppm – 252.61 ppm of which 23% samples are very low, 26% samples are low, 6% samples are medium, 10% samples are optimum, 3% samples are high and 32% samples are very high in fertility status. The sulfur content of soil samples of Rajshahi division ranges from 0.23 ppm – 16.28 ppm of which 82% samples are very low, 9 % samples are low and 9% samples are medium in fertility status. The sulfur contents of soil samples of Khulna division ranges from 0.03 ppm – 258.09 ppm where 92% samples are very low and 8% samples are very high in fertility status. The sulfur content of soil samples of Barishal division ranges from 0.84 ppm – 43.39 ppm of which 33% samples are very low, 25% samples are low, 17% samples are medium, 17% samples are high and 8% samples are very high in fertility status. The sulfur content of soil samples of Rangpur division ranges from 0.64 ppm – 68.61ppm of which 50% samples are very low, 28% samples are low, 7% samples are high and 15% samples are very high. The sulfur contents of soil samples of Sylhet division ranges from 6.83 ppm – 9.82 ppm of which 50% samples are very low and 50% samples are low and the sulfur content of soil samples of Mymensingh division ranges from 1.47 ppm – 30.73 ppm of which 50% samples are very low, 20% samples are low, 10% samples are medium, 10% samples are optimum and 10% samples are high in fertility status.

Zinc

The zinc content of soil samples of Dhaka division ranges from 0.64 ppm– 37.76 ppm of which 9% samples are low, 14% samples are medium, 7% samples are optimum, 9% samples are high and 61% samples are very high in fertility status. The zinc content of soil samples of Chattogram division ranges from 0.68ppm – 2.65 ppm of which 42% samples are low, 32% samples are medium, 23% samples are optimum and 3% samples are very high. The zinc content of soil

samples of Rajshahi division ranges from 1.16 ppm – 8.52 ppm of which 9% samples are medium, 27% samples are optimum, 27% samples are high and 37 % samples are very high. The zinc content of soil samples of Khulna division ranges from 0.68 ppm – 9.2 ppm of which 31% samples are low, 23% samples are medium, 15% samples are optimum, 8% samples are high and 23% samples are very high. The zinc content of soil samples of Barishal division ranges from 0.74 ppm – 3.77 ppm of which 17% samples are low, 25% samples are medium, 17% samples are optimum, 16% samples are high and 25% samples are very high. The zinc content of soil samples of Rangpur division ranges from 1.02ppm – 3.62 ppm of which 43% samples are medium, 21% samples are optimum, 27% samples are high and and 37% samples are very high. The zinc contents of soil samples of Sylhet division ranges from 0.75 ppm – 0.84 ppm of which 100% samples are low and the zinc content of 10 soil samples of Mymensingh division ranges from 0.88 ppm – 12.85 ppm of which 10% samples are low, 10% samples are medium, 50% samples are optimum and 10% samples are high and 20% samples are very high in fertility status.

Boron

The boron content of 44 soil samples of Dhaka division ranges from 0.003 ppm– 1.70 ppm of which 50% samples are very low, 11% samples are low, 5% samples are medium, 7% samples are optimum, 9% samples are high and 18% samples are very high in fertility status. The boron content of 31 soil samples of Chattogram division ranges from 0.001 ppm – 0.89 ppm of which 61% samples are very low, 13% samples are low, 10% samples are medium, 13% samples are optimum and 3% samples are very high. The boron content of soil samples of Rajshahi division ranges from 0.12 ppm – 0.85 ppm of which 19% samples are very low, 27% samples are low, 27% samples are medium, 9% samples are optimum, 9% samples are high and 9% samples are very high. The boron content of soil samples of Khulna division ranges from 0.003 ppm – 0.58 ppm of which 39% samples are very low, 15% samples are low, 31% samples are medium, and 15% samples are optimum. The boron content of soil samples of Barishal division ranges from 0.11 ppm – 1.48 ppm of which 8% samples are very low, 17% samples are low, 17% samples are medium, 17% samples are optimum and 41% samples are very high. The boron content of soil samples of Rangpur division ranges from 0.001 ppm – 1.40 ppm of which 57% samples are very low, 22% samples are low, 7% samples are optimum and 14% samples are very high. The boron

content of soil samples of Sylhet division ranges from 0.48 ppm – 0.74 ppm of which 50% samples are optimum and 50% samples are high and the boron content of 10 soil samples of Mymensingh division ranges from 0.11 ppm – 0.78 ppm of which 20% samples are very low, 20% samples are low, 20% samples are medium, 30% samples are high and 10% samples are high in fertility status.

1.4 Analyses of soil samples under Land and Soil Resources Utilization Guide Updating Program

Preparation of Land and Soil Resources Utilization Guide is one of the most important programs of SRDI. SRDI created huge database on soil physical and chemical properties through the program. A total of 68 soil samples of SauriaUpazilla under Manikgonj district have been analyzed in the year 2017-2018. Different macro and micro nutrient elements like nitrogen, phosphorus, potassium, sulfur, boron zinc, calcium, magnesium, copper, iron, manganese as well as pH and organic matter content of the soil were determined in the laboratory following the standard procedures of respective elements.

1.5 Farmers service through Mobile Soil Testing Laboratory

Mobile Soil Testing Laboratory (MSTL) is one of the popular programs of SRDI. The objective of the service is to ensure the application of balanced fertilizer doses to the farmers' fields which will create awareness among the farmers on the benefits of soil test based balanced fertilizer application. In the 2017-2018 fiscal year MSTL Jamuna, operated by Central Laboratory, provided service to 756 farmers of 16 upazillas of 8 districts in two seasons (Rabi and Kharif) under this program. In the Rabi season Manikganj Sadar (Manikganj), Delduar (Tangail), Basail (Tangail), Mymensing Sadar (Mymensing), Barhatta (Netrokona), Nikli (Kishoregonj), Raipura (Narsingdi), Nawabganj (Dhaka) and in the Kharif season Sripur (Gajipur), Daulatpur(Manikgonj), Tangail Sadar (Tangail), Bhuapur (Tangail), Purbadhala (Netrokona), Nandail (Mymensing), Katiyadi (Kishoregonj), Palash (Narsingdi) Upazilas were covered under this program.

1.6 Quality of fertilizers in 2017-2018 fiscal year under quality control activities

In 2017-2018 fiscal year Central Laboratory of SRDI received 1025 Fertilizer samples from different government and private sectors and analyzed all fertilizer samples for quality assurance. Out of that fertilizer samples cent percent Urea, Di-ammonium phosphate (DAP), Sulphate of Potash (SOP) and Ammonium sulfate were standard sample as per government specification. Liming material Dolomite was also 100% standard as per government specification. Central Laboratory received 47 TSP samples. Out of 47 TSP fertilizers only 2 samples were adulterated that is 96% TSP was standard. In case of Murate of Potash (MoP) 98%, Gypsum 96%, Magnesium sulfate 99% samples were standard. On the other hand, 57% NPKS mixed fertilizer were found adulterated at different levels of impurities. In case of micronutrient fertilizer the maximum adulteration was observed in Zinc (Zn) fertilizers. In 2017-2018 fiscal year 158 Zinc sulfate fertilizer samples were analyzed some in the form of Zinc sulfate monohydrate and zinc sulfate heptahydrate. Out of 158 Zinc sulfate fertilizer samples only 35 samples were standard which was 28% of the received Zinc sulfate fertilizer samples and rest 78% was adulterated at different degrees of adulteration. The adulteration of Zinc fertilizers was not only concentrated in the deficiency of Zn content but also the excessive presence of heavy metal contamination. Another government specified important source of Zinc is Chelated Zinc. Thirty three (33) Chelated zinc fertilizer samples have been analyzed in this year by Central Laboratory where 28 samples of Chelated Zinc were found standard (85%). Other widely used micronutrient fertilizer is Boron fertilizer. Boron is the most important micro nutrient fertilizer for crop production. Boric acid and Solubor are the familiar fertilizer forms. In the last fiscal year Central Laboratory analyzed 69 Solubor Boron where 67 samples were standard which was 97% of the total sample. On the other hand, it analyzed 39 Boric acid fertilizer samples where 36 samples were standard which was 92% of the total sample. In the last year we analyzed 138 organic fertilizer samples where 49% organic fertilizers were standard. It was found that the percent of adulteration in the physically blended NPKS mixed fertilizers was 57% (Table 1). It was also reported from Department of agriculture extension (DAE) that due the adverse effect of adulterated NPKS mixed fertilizer on crop production farmers are not interested in using this fertilizer. It is notable that the quality control activities of this organization has been playing an important role reducing the percent of adulteration remarkably as the adulteration in fertilizer was much higher in the previous time.

Table 2. Quality of different fertilizers analyzed by Central Lab in 2017-2018

Fertilizer	Total	Standard	Adulterated	Standard (%)	
				Quality	Adulterated
Urea	48	48	0	100	0
TSP	47	45	2	96	4
DAP	50	50	0	100	0
MOP	58	57	1	98	2
SOP	45	45	0	100	0
Gypsum	73	70	3	96	4
Zinc Sulphate	158	35	123	22	78
Chelated Zinc	33	28	5	85	15
Mg Sulfate	174	173	1	99	1
Boric Acid	39	36	3	92	8
Solubor Boron	69	67	2	97	3
Ammonium Sulfate	56	56	0	100	0
Dolomite	12	12	0	100	0
Organic	138	67	71	49	51
NPKS	14	6	8	43	57
Others	11	6	5	55	45
Total	1025	801	224		

*Others fertilizers included Silvagin, Silvamix, Nutraphos-N, Nutraphos-24, Copper Vit, Wuxl super, Peak, Biochar, American NPK.

1.6.1 Trend of micro nutrient fertilizer quality

Zinc fertilizer

In the last four consecutive fiscal years zinc (Zn) fertilizer quality was not improved rather to some extent its quality is decreased (Fig. 1). In 2014-2015 fiscal years the adulterated percent of Zn fertilizers were 72%. On the other hand, in the last 2017-2018 fiscal years the adulterated percent of Zn was 78%. This important micro nutrient fertilizer is produced, imported and marketed by private sector. Some unfair traders import low quality Zn fertilizers from abroad. Some manufacturer's produce this fertilizes using below standard raw materials. More over these fertilizers are adulterated not only concentrated in the deficiency of nutrient content (Zinc and Sulfur) but also the excessive presence of heavy metals like cadmium, lead, nickel, chromium etc. To ensure the quality of Zn fertilizer at field level monitoring should be strengthened through the fertilizer inspectors of DAE and other law enforcing agencies. Generally fertilizer inspectors of DAE are sampling from fertilizer dealers and sub-dealers. In case of Zn fertilizers it would be more effective if samples are collected lot by lot from manufacturing factories.




Figure 22. Trend of Zn fertilizers quality last four consecutive fiscal years




Figure 23. Trend of Chelated-Zn fertilizers quality last four consecutive fiscal years

Chelated-Zn fertilizers

Chelated zinc fertilizer is an expensive foliar applicable fertilizer which easily available to plants and also important source of Zn for crops. In the last four consecutive fiscal years Chelated Zinc fertilizer quality was improved. In the 2014-2015 fiscal years the standard Chelated Zinc fertilizer percent was 74%. On the other hand, in the last 2017-2018 fiscal years the percent of quality Chelated zinc fertilizer was 85% (Fig. 2).




Figure 24. Trend of Solubor quality last four consecutive fiscal years




Figure 25. Trend of Boric acid quality last four consecutive fiscal years

Solubor and Boric acid

There are four specified Boron fertilizers in Bangladesh viz. Solubor, Boric acid, Fertibor and Granubor. Among the four fertilizers Solubor and Boric acid are most familiar to the farmers of country. In the last four consecutive fiscal years Solubor fertilizer quality was improved. In the 2014-2015 fiscal years the standard Solubor fertilizer percent was 92%. On the other hand, in the last 2017-2018 fiscal years the percent of quality Solubor fertilizer was 97% (Fig. 3). In the last four consecutive fiscal years Boric acid fertilizer quality was improved remarkably. In the 2014-

2015 fiscal years the standard Boric acid fertilizer percent was 73%. On the other hand, in the last 2017-2018 fiscal years the percent of quality Boric acid fertilizer was 92% (Fig. 4).




Figure 26. Trend of Organic fertilizer quality last four consecutive fiscal years

1.6.2 Trend of organic fertilizer quality

Government specified organic fertilizer in April, 2008 with inclusion of specific physical and chemical parameters. In 2014-2015 fiscal year 74% organic fertilizers were substandard /adulterated where as in 2017-2018 the adulteration/ substandard organic fertilizers were 51% (Fig. 5). It indicates the trend of quality improvement of organic fertilizers. The improvement of organic fertilizers quality might be intensive monitoring and awareness of the organic fertilizer producers. Many young and educated entrepreneurs are involved in producing and marketing organic fertilizers by following government approved specification during the recent past. Organic fertilizers were substandard /adulterated in various aspects including nutrient deficiency, low organic carbon, presence of excessive moisture, absence of desirable C:N ratio and presence of excessive pollutant like heavy metal content like lead (Pb), Cadmium (Cd), Chromium (Cr) and Nickel (Ni) etc.

References

- BARC, 2005. Fertilizer recommendation guide. Bangladesh Agricultural Research Council (BARC), Farmgate, New airport road, Dhaka-1215.
- Bray, R. H. and L. T. Kurtz. 1945. Determination of total, organic, and available forms of phosphorus in soils. Soil Sci. 59: 39-45.

- Bremner, J. M. and C. S. Mulvaney. 1982. Total nitrogen. In *Methods of soil analysis, Part 2, Chemical and microbiological properties*, (e.d.) A. L. Page, R. H. Miller, and D. R. Keeney. *Soil Sci. Soc. Am. J.* 595-624.
- Barker, D. E. and N. H. Surh. 1982. Atomic absorption and flame emission spectroscopy. In *Methods of soil analysis, Part 2, Chemical and microbiological properties*, (e.d.) A.L. Page, R.H. Miller, and D.R. Keeney. *Am. Soc. of Agron. and Soil Sci. Soc. Am. J.* 13-26.
- Hasan, M. N., and Rahman, M. L., 2015. A Comparative Assessment of Soil Fertility in Birganj Upazilla of Dinajpur District and Hatibandha Upazilla of Lalmonirhat District of Bangladesh, 53 pp.
- Kundu, S., R. Bhattacharyya, V. Prakash, B. N. Ghosh and H. S. Gupta. 2006. Carbon sequestration and relationship between carbon addition and storage under rain fed soybean-wheat rotation in a sandy loam soil of the Indian Himalayas. *Soil Tillage Res.* 92: 87-95.
- Lal, R. 2006. Enhancing crop yields in the developing countries through restoration of the soil organic carbon pool in agricultural lands. *Land Degrad. Develop.* 17: 197-209.
- Lindsay, W. L. and W. A. Norvell. 1978. Development of a DTPA soil test for zinc, iron, manganese and copper. *Soil Sci. Soc. Am. J.* 42: 421-428.
- McLean, E. O. 1982. Soil pH and lime requirement. In *Methods of soil analysis, Part 2, Chemical and microbiological properties*, (e.d.) A. L. Page, R. H. Miller and D. R. Keeney. *Am. Soc. Agron.* pp 199-224.
- Mikhailova, E. A. and C. J. Post. 2006. Effects of land use on soil inorganic carbon stocks in the Russian Chernozem. *J. Environ. Qual.* 35: 1384-1388.
- Nelson, D. W. and L. E. Sommers, 1982. Total carbon, organic carbon, and organic matter. In: *Methods of soil analysis, Part 2, Chemical and microbiological properties*, A. L. Page, R. H. Miller and D. R. Keeney (eds). *Am. Soc. Agron.* 539-579.
- Page, L. R., R. H. Mitter and D.R. Keeney. 1989. *Methods of Soil Analysis. Part II*, and ed. American Soc. Agron. Inc. Pub. Madison Wisconsin, USA.
- Tallahassee, 2013. Statistix 10, Analytical software, Florida, USA.
- Wolf, B. 1974. *Soil Sci. plant Anal.* 5:39-44.

Proposed research programs for 2018-2019

1. Study on Changing Pattern of Soil Fertility of Dhamri Upazila, Dhaka; Saturia Upazila, Manikganj and Kalihati Upazila, Tangail District under Changing Climate

Introduction

A common belief is that soil productivity is decreasing day by day with increasing cropping intensity and imbalanced use of fertilizers and lack of proper soil and crop management practices. This is particularly true for intensive crop cultivated areas of Bangladesh. Crop productivity of Bangladesh seems to be high and improving, cropping intensity is above national average but due to imbalance use of fertilizers and high cropping intensity, soil acidity is increasing while base materials are decreasing from the top soils. During the last 30 years the agricultural sector of the country has developed significantly, from traditional to intensive system, production is also much higher than the previous years; poverty level decreased remarkably and all other development indicators are in positive trend. However, there has been pressure on natural resources, particularly on soil fertility. Degradation of soil quality in terms of losing its base material content is a serious threat in this aspect. So, to overcome these problems it is important to adopt conservation farming or climate smart farming for sustainability. The reality is that for long time farmers used urea indiscriminately for production without considering balanced use of other essential nutrient like phosphorus (P), potassium (K), sulfur (S) and micronutrients. It was observed that in a short period of time 23% organic matter depletion has been found in the northern region which is very alarming and both Calcium and Magnesium content is also considerably lowering down (Hasan and Rahman, 2015). It is apprehended that the above soil fertility scenario is almost same all over the country. Farmers are using lime to improve soil pH and nutrient availability but the rate of application is not yet set. So they do not get the actual benefit from liming and fertilizer use. The term sustainability and production system are very important as there is a decreasing sign of fertilizer response. Considering the above perspective Dhamri Upazila of Dhaka district, Saturia Upazila of Manikganj district and Kalihati Upazila of Tangail District were selected for the study with the objectives to determine the changes in soil quality in terms of available nutrient content, changes in organic matter content and soil pH over time period.

Methods and Materials

To carry out the study the Upazila named Dhamri of Dhaka district, Saturia of Manikganj district and Kalihati of Tangail district are selected. Dhamri Upazila is under the Agro-ecological zone (AEZ) Active Bhrahmaputra-Jamuna Floodplain, Young Bhrahmaputra-Jamuna Floodplain, Old Brahmaputra Floodplain and mixed Active Bhrahmaputra-Jamuna Floodplain and Madhupur Tract located in the middle part of Bangladesh. Soil Resource Development Institute (SRDI) conducted semi-detailed soil survey at Dhamri Upazila in 2002 and collected 89 soil samples from different land types and soil series. Those chemical databases of 2002 are considered as baseline for the present study. Under updating program of Upazila Land and Soil Resource Utilization Guide popularly known as Thana Nirdeshika, SRDI also collected 133 soil samples from different land type, soil series and AEZs in 2018. A comparative study will be done on nutrient content, soil reaction, soil organic matter content considering various aspect like land type, soil series, AEZs. Saturia Upazilla is under the AEZ Young Bhrahmaputra-Jamuna Floodplain and Active Bhrahmaputra-Jamuna Floodplain located in the middle part of Bangladesh. Semi detail survey of Saturia Upazila of Manikganj district was done 1999. During that period 48 soil samples from Saturia Upazila and analyzed in the laboratory for chemical parameters. Those data base will be considered as base line database for the present study. Under updating program of Thana Nirdeshika, SRDI also collected 68 soil samples from different land type, soil series and AEZs in 2018. A comparative study will be done on nutrient content, soil reaction, soil organic matter content considering various aspect like land type, soil series, AEZs. Kalihati Upazilla is under the AEZ Young Bhrahmaputra-Jamuna Floodplain and Active Bhrahmaputra-Jamuna Floodplain and Madhupur Tract also located in the middle part of Bangladesh. Semi detail survey of Saturia Upazilla of Manikganj district was done 1998. During that period 124 soil samples from Kalihati Upazilla and analyzed in the laboratory for chemical parameters. Those data base will be considered as base line database for the present study. Under updating program of Thana Nirdeshika, SRDI also collected 106 soil samples from different land type, soil series and AEZs in 2018. A comparative study will be performed for nutrient content, soil reaction, soil organic matter content considering various aspect like land type, soil series, AEZs. Land use, use of organic matter, crop yield and fertilizer use related information was also collected during the survey. Chemical analysis will be done in the Central Laboratory of Soil Resource Development Institute (SRDI) by following worldwide standard methods. Soil pH was

determined by Glass Electrode pH meter method with soil water ratio 1:2.5 (McLean, 1982), Organic matter by Walkley-Black method (Nelson and Sommers, 1982), total N by Kjeldhal system (Bremner and Mulvaney, 1982), available P by Bray and Kurtz Method (1945), K by ammonium acetate extraction method (Barker and Surh, 1982), available S was determined by Turbidimetric method (Page *et al.*, 1989), available Cu, Fe, Mn and Zn was determined by DTPA Extraction method (Lindsay and Norvell, 1978.) and B is determined by Calcium chloride extraction method (Wolf, 1974).

2. Assessment of Chemical and physical Properties of Composting Raw Materials

Introduction

Soils play an important role in the global carbon budget and therefore, it is important to account for soil carbon stocks in different soils under different land use systems (Mikhailova and Post, 2006). Sustaining soil organic matter (SOM) is of paramount importance with respect to availability of plant nutrients and improvement of the soil's physical, chemical and biological properties (Kundo *et al.*, 2006). Maintenance of soil organic carbon (SOC), a major component of SOM, is essential for the sustainable agricultural production as declining SOC generally leads to decreased crop productivity (Lal, 2006). Soils of Bangladesh have low reserves of carbon and plant nutrients due to increasing cropping intensity, higher rates of decomposition of organic matter under prevailing hot and humid climate, use of lesser quantity of organic manure, little or no use of green manure. According to BARC (2005), the highest depletion of soil carbon has been reported in soils of Meghna river floodplain (35%), followed by Madhupur Tracts (29%), Brahmaputra floodplain (21%), Old Himalayan piedmont plain (18%) and Ganges floodplain (15%). Carbon stocks are not only critical for the soil to maintain its productivity and environmental functions but also play an important role in the global carbon cycle. Soil organic matter is essential to improve soil quality, increase agronomic productivity and use efficiency of inputs like fertilizers and water thus helps maintain or restore the capacity of soil to perform its production and environmental function on a sustainable basis.

Government specified organic fertilizer in April, 2008 with inclusion of so many physical and chemical features. In 2014-2015 74% organic fertilizers were adulterated where as in 2017-2018 the adulteration percent of organic fertilizers reduced to 51%. Many young and educated

entrepreneurs are coming forward in producing and marketing organic fertilizers by following government approved specification during the recent past. Organic fertilizers were adulterated in various aspects including nutrient deficiency, low organic carbon, presence of excessive moisture, absence of desirable C:N ratio and presence of excessive pollutant like heavy metal lead (Pb), Cadmium (Cd), Chromium (Cr) and Nickel (Ni) etc. In maximum case the adulteration happened due to lack of adequate knowledge and information about the quality of composting materials and composting techniques. Maximum people those are involved in producing organic fertilizers do not know carbon content, C:N ratio, others chemical properties of the composting materials. So organic fertilizers produced by them can not comply the government approved specification. Whereas use of quality organic fertilizers beside of chemical fertilizers essential to maintain the soil health. Soil organic matter is a prime concern to maintain soil fertility and increase the use efficiency of chemical fertilizers and other agricultural inputs specially irrigation. More over in recent past roof gardening or roof agriculture is becoming popular in urban areas of Bangladesh which requires huge quantity of quality organic fertilizers. Therefore it is an urgent issue to study the potentiality of different composting materials for the production of good quality organic fertilizer. To consider the above perspective the study is taken to identify the potentiality of different organic sources for the production of quality organic fertilizers.

Methods and Materials

To conduct the study different composting materials like Water hyacinth, Poultry waste, Kitchen waste, Coconut coir, Press mud, Tobacco dust, Vegetable waste, Cowdung, Pine apple leaves, and different Oil cake like oil cake, Soya cake, Neem cake, Mahagani cake etc. will be collected from three locations. Each composting material will be collected from three different locations. From each location three samples of each composting materials will be collected. Chemical properties like pH, organic carbon, total nitrogen, C:N ratio, phosphorus, potassium, sulfur, zinc, copper, cadmium, lead, nickel, chromium will be determined. Chemical and physical properties of these composting materials will be determined following standard methods approved the Government of Bangladesh. Statistix 10 software will be used to analyze the data. ANOVA and univariate analysis were performed. Means were separated by LSD at 5% level of significance.

3. Effect of Irrigation Water and Fertilizer Use on Soil pH and Organic Matter in the Soils of Barind Tract

Introduction

The Barind Tract is an elevated Pleistocene Terraces (about 11-48 m MSL) in northwestern Bangladesh. The geoscientists thought that the tract may have been evolved due to tectonic upliftment and /or exists as an erosional geomorphic feature but nobody agrees the same opinion. This problem represents different questions with variable dimension in the groundwater problem in the area. Most of the areas where underlying Madhupur clay have been uplifted and cut into deep valleys. The topsoil is grey silt loam, is strongly puddle and has compact ploughpan at the base. The subsoil is grey, brightly mottled yellow-brown, silt loam to clay and is very porous. Deep grey Terrace soils and grey valley soils are the major components of the general soil types of the area. General fertility status is low to medium. Status of organic matter was very low to low; fertility status of N, P and K was very low to low and fertility status of Zn and B was low to medium. Top soil pH was ranged from very strongly acidic to slightly alkaline. In recent past it was noticed to the agricultural scientist that soil pH in some soils of Barind Tract have been changed. Before implementation of Barendra Multipurpose Irrigation project cropping intensity was less than national average. But after the implementation of the irrigation project cropping intensity has been increased and huge underground water is harvested for irrigation. Water management and fertilizer use may alter the soil pH, organic matter status of the area. Therefore, urgent study is necessary to determine the changing pattern of soil pH, organic matter content and other nutrient content under the altered agricultural practice.

Methods and Materials

To carry out the study three Upazilas of Barind Tract under the command area of Barindra Multipurpose Irrigation project will be selected where irrigated and non- irrigated farming systems exit. From each Upazila 50 soil samples from irrigated area and 50 soil samples from non-irrigated area will be collected for the study. Underground irrigation water quality will be determined to examine the effect irrigation water on soil properties. Chemical analysis will be done in the Central Laboratory of Soil Resource Development Institute (SRDI) followed by worldwide standard methods. Soil pH will be determined by Glass Electrode pH meter method


with soil water ratio 1:2.5 (McLean, 1982), Organic matter by Walkley-Black method (Nelson and Sommers, 1982), total N by Kjeldhal system (Bremner and Mulvaney, 1982), available P by Bray and Kurtz Method (1945), K by ammonium acetate extraction method (Barker and Surh, 1982), available S was determined by Turbidimetric method (Page *et al.*, 1989), available Cu, Fe, Mn and Zn was determined by DTPA Extraction method (Lindsay and Norvell, 1978.) and B is determined by Calcium chloride extraction method (Wolf, 1974). Statistix 10 software (Tallahassee, 2013) was used to analyze the data. ANOVA and univariate analysis were performed. Means were separated by LSD at 5% level of significance.

2.9 Upazila Nirdeshika Cell

Publication of Updated Nirdeshika from revenue budget

Two hundred and twenty seven reports of Upazila Nirdeshika has been updated till June 2018 with a series of coordinated effort such as semi-detailed soil survey for updating of relevant maps, collection of land quality and soil characteristics data, soil sample collection for laboratory analysis followed by report writing with the processing of relevant updated data generated during field survey and laboratory analysis. SRDI scientists engaged in district offices usually conduct the soil survey and prepare draft report for respective surveyed Upazila. Editing of the report is done initially by regional head (Principal Scientific Officer) who supervise survey based soil mapping and finally by the editorial board acting particularly as the approval authority of Upazila Niredeshika publication. The activities involved in the Upazila Nirdeshika updating system can be expressed as a flow-chart.

The activities of updating Nirdeshika previously were funded solely by SRDI projects (till June 2016) and the activities linked with the Nirdeshika updating have also been conducted utilizing revenue budget from the financial year 2016–2017. The entire activity is coordinated by Upazila Nirdeshika Cell at SRDI Head Office. Thirty updated Nirdeshika has been published during the period of 2017-2018 under revenue budget of SRDI (Table 1).



Flow chart on activities of Upazila Nirdeshika Cell

Table 3. List of Updated Upazila Nirdeshika published by expending the revenue budget during 2017-2018

No.	Name of Upazila (Nirdehika updated)	District	Year of 1 st Edition of Nirdeshika	Area of the Upazila (ha)
1	Muladi	Barishal	1999	23,300
2	Durgapur	Rajshahi	1999	19,504
3	Taltoli	Barguna	1997	24,242
4	Shoilkupa	Jhenaidah	1993	37,345
5	Kumarkhali	Kushtia	1995	25,838
6	Burichang	Cumilla	2001	16,377
7	Phulpur	Mymensingh	2000	28,469
8	Tungibari	Munshiganj	1991	11,933
9	Gabtali	Bogura	1989	23,971
10	ChapaiNawabganjSadar	ChapaiNawabganj	1996	40,852
11	Kaptai	Rangamati	1997	27,324
12	Birganj	Dinajpur	1999	41,312

No.	Name of Upazila (Nirdehika updated)	District	Year of 1 st Edition of Nirdeshika	Area of the Upazila (ha)
13	Anowara	Chattogram	1996	15,869
14	Naniarchar	Rangamati	1999	38,793
15	Hossainpur	Kishoregonj	1997	11,972
16	ChandpurSadar	Chandpur	1999	27,714
17	keshobpur	Jessore	1997	25,853
18	BholaSadar	Bhola	1993	41,314
19	Shonagazi	Feni	1999	23,564
20	Uzirpur	Barishal	2000	24,841
21	Bijoy Nagar	Brahmanbaria	1993	21,570
22	Alphadanga	Faridpur	2000	13,601
23	Phulbaria	Mymensingh	1992	39,870
24	Nagorpur	Tangail	1995	26,720
25	Madhupur	Tangail	2001	47,772
26	Bahubol	Habiganj	2002	24,861
27	Kutubdia	Cox` s bazar	1999	15102
28	Sribordi	Sherpur	1992	25,996
29	Chatak	Sylhet	1999	44,441
30	Shaturia	Manikganj	1999	13,846

Chapter 3: Activities of Field Offices

3.1. Updating Upazila Land and Soil Resource Utilization Guide (Upazila Nirdeshika) through Semi-detailed Soil Survey

(a) Introduction: Upazila Land and Soil Resource Utilization Guide (Upazila Nirdeshika) is being used as a tool for agricultural development activities at Upazila level. Beginning from 1986 all the 459 Upazila Nirdeshika publication was completed by June 2002. With passing of time changes in edaphic properties, land use and fertility status of soil are being observed along with physical changes due to population growth, rapid industrialization and urbanization, deforestation, unplanned construction of roads, culverts, dams, blocking of upstream flow of rivers, indiscriminate use of chemical pesticides & fertilizers, global warming etc. These changed scenarios need to be incorporated in the Nirdeshika. That is why it becomes imperative to update the existing Upazilla Nirdeshika to enrich the database and make the Nirdeshika more useful to the field level users with the following objectives.

(b) Objectives

- To update the land, soil and land use database for local and regional level agricultural development planning.
- To update the soil fertility database.
- To accommodate the changes due to infrastructure developments (roads, homestead, embankments etc.).
- To make it user friendly and include executive summary of the whole text for the decision makers.

(c) Methodology

Base Materials: Existing Upazila Soil and Landform Map (1:50,000), aerial photographs of approximate scale of 1:25,000 of 1:30,000, topographic maps (1:50,000), DLR maps (1: 63,360) are used as field base maps.

Methods: Based on interpretation of recent aerial photos a photo interpretative Soil and Landform Map is prepared with the help of existing one. The map consists of legend depicting soil mapping unit(s), land type, Mrittika Dal (Soil group).

A planned ground truthing is made by checking catena wise soil, mapping unit boundary following regular traverse and grid as needed in semi-detailed survey.

Soils are examined as often as necessary along traverse lines. For each 200 hectares of land, one soil sample is collected on the basis of land type and Mrittika Dal. The sampling intensity is increased as and when necessary, according to the complexity of Mrittika Dal.

Composite soil samples are collected from adjacent to or possibly nearer point of previous sampling sites with GPS reading so as to compare the changes of nutrient status due to intensive

cultivation of modern varieties of different crops with imbalanced application of chemical fertilizers and climate changes.

Mini pits are described by digging shallow pits (where necessary) along each traverse. Soil samples are taken in correlation boxes (if necessary) from Mrittika Dal for correlation.

During soil sample collection, information on inundation depth, cropping pattern, constraints for agricultural development etc. are collected from field observation and farmers interview.

Collected composite soil samples are analyzed in the laboratory and updated Upazila Nirdeshika is prepared incorporating land, soil resource and fertility data.

Table 1: Progress of Upazila Nirdeshika Updating

District	Field Survey		Map finalization		Draft Report Preparation		Final Report Preparation	
	Target	Achievement	Target	Achievement	Target	Achievement	Target	Achievement
Regional Office, Rajshahi								
Rajshahi	Naldunga	Naldunga	Naldunga	Naldunga	Dhamoirhat	Dhamoirhat	Godagari	Godagari
	Gomostapur	Gomostapur	Gomostapur	Gomostapur	Badalgachi	Badalgachi	Chapainawabganj Sadar	Chapainawabganj Sadar
	Porsha	Porsha	Porsha	Porsha	-	-	Durgapur	Durgapur
Dinajpur	Fulbari	Fulbari	Fulbari	Fulbari	Khansama	Khansama	Birganj	Birganj
	Pirganj	Pirganj	Pirganj	Pirganj	Chirirbandar	Chirirbandar	Kaharol	Kaharol
	Debiganj	Debiganj	Debiganj	Debiganj	-	-	-	-
Bogura	Shajahanpur	Shajahanpur	Shajahanpur	Shajahanpur	Nandigram	Nandigram	Naogaon Sadar	Naogaon Sadar
	Panchbibi	Panchbibi	Panchbibi	Panchbibi	Shariakandi	Shariakandi	Gaboli	Gaboli
	-	-	-	-	-	-	Dhunat	Dhunat
Rangpur	Jaldhaka	Jaldhaka	Jaldhaka	Jaldhaka	Gaibandha Sadar	Gaibandha Sadar	Fulbari	Fulbari
	Nageshwari	Nageshwari	Nageshwari	Nageshwari	Rajibpur	Rajibpur	Kurigram Sadar	Kurigram Sadar
	Lalmonirhat	Lalmonirhat	Lalmonirhat	Lalmonirhat	-	-	-	-
Pabna	Raiganj	Raiganj	Raiganj	Raiganj	Belkuchi	Belkuchi	Lalpur	Lalpur
	Kazipur	Kazipur	Kazipur	Kazipur	-	-	-	-
Chattogram Regional Office, Cumilla								
Cumilla	Homna	Homna	Sadar Dakshin	Sadar Dakshin	Sadar Dakshin	Sadar Dakshin	Burichong	Burichong
	Bancharampur	Bancharampur	Kachua	Kachua	Kachua	Kachua	Chandpur Sadar	Chandpur Sadar
	-	-	Shahrasti	Shahrasti	Shahrasti	Shahrasti	Bijoynagar	Bijoynagar
Noakhali	Noakhali Sadar	Noakhali Sadar	-	-	-	-	Sonagazi	Sonagazi
Chattogram	Lohagara	Lohagara	Satkania	Satkania	Satkania	Satkania	Anwara	Anwara
	-	-	Ukhiya	Ukhiya	Ukhiya	Ukhiya	Pekua	Pekua
	-	-	-	-	-	-	Kutubdia	Kutubdia
Rangamati	Rajasthali	Rajasthali	Ramgarh	Ramgarh	Ramgarh	Ramgarh	Kaptai	Kaptai
	-	-	-	-	-	-	Naniarchar	Naniarchar
Regional Office, Sylhet								
Sylhet	Dakshin Sunamganj	Dakshin Sunamganj	-	-	-	-	Chatok	Chatok
Moulvibazar	Kamalganj	Kamalganj	-	-	-	-	Bahubal	Bahubal
	Nobiganj	Nobiganj	-	-	-	-	-	-
Regional Office, Dhaka								
Dhaka	Kaliganj	Kaliganj	Kaliganj	Kaliganj	Dhamrai	Dhamrai	Tongibari	Tongibari
	Shirajdekhan	Shirajdekhan	Shirajdekhan	Shirajdekhan	Kaliakoir	Kaliakoir	-	-
	Nowabganj	Nowabganj	Nowabganj	Nowabganj	Rupganj	Rupganj	-	-
Tangail	Shibaloy	Shibaloy	Shibaloy	Shibaloy	Nagarapur	Nagarapur	Sribordi	Sribordi
	Gopalpur	Gopalpur	Gopalpur	Gopalpur	Shibganj	Shibganj	Dhunot	Dhunot
	-	-	-	-	Kalihati	Kalihati	Dhanbari	Dhanbari
	-	-	-	-	Islampur	Islampur	Bhuapur	Bhuapur

Faridpur	Kalkini	Kalkini	Kalkini	Kalkini	Kashiani	Kashiani	Muladi	Muladi
	Kotalipara	Kotalipara	Kotalipara	Kotalipara	Bhanga	Bhanga	Kalukhali	Kalukhali
	Naria	Naria	Naria	Naria	-	-	-	-
Jamalpur	Madan	Madan	Madan	Madan	Dewanganj	Dewanganj	-	-
	-	-	-	-	Nagarpur	Nagarpur	-	-
Mymensingh	Bajitpur	Bajitpur	Bajitpur	Bajitpur	Atpara	Atpara	Tarakanda	Tarakanda
	Kotiadi	Kotiadi	Kotiadi	Kotiadi	Gaforgaon	Gaforgaon	Fulbaria	Fulbaria
	Bhaluka	Bhaluka	Bhaluka	Bhaluka	Kuliarchan	Kuliarchan	Fulpur	Fulpur
	-	-	-	-	Mithamoin	Mithamoin	Hossainpur	Hossainpur

District	Field Survey		Map finalization		Draft Report Preparation		Final Report Preparation	
	Target	Achievement	Target	Achievement	Target	Achievement	Target	Achievement
Regional Office, Khulna								
Khulna	Rampal	Rampal	Rampal	Rampal	Tala	Tala	-	-
	Kachua	Kachua	Kachua	Kachua	Paikgacha	Paikgacha	-	-
	Dacope	Dacope	Dacope	Dacope	-	-	-	-
Jashore	Kaligong	Kaligong	Kaligong	Kaligong	Monirampur	Monirampur	Keshobpur	Keshobpur
	Shalikhha	Shalikhha	Shalikhha	Shalikhha	Debhata	Debhata	Kalaroa	Kalaroa
	Mongla	Mongla	Mongla	Mongla	-	-	-	-
	Saronkhola	Saronkhola	Saronkhola	Saronkhola	-	-	-	-
	Morelgonj	Morelgonj	Morelgonj	Morelgonj	-	-	-	-
Kushtia	Horinakundu	Horinakundu	Horinakundu	Horinakundu	-	-	-	-
	-	-	-	-	-	-	Alamdanga	Alamdanga
	-	-	-	-	-	-	Kumarkhali	Kumarkhali
Regional Office, Barishal								
Barishal	Nazirpur	Nazirpur	Nazirpur	Nazirpur	Bhola Sadar	Bhola Sadar	Wazirpur	Wazirpur
	Charfassion	Charfassion	Charfassion	Charfassion	-	-	Muladi	Muladi
	Manpura	Manpura	Manpura	Manpura	-	-	-	-
Patuakhali	Kathalia	Kathalia	Kathalia	Kathalia	Betagi	Betagi	Barguna Sadar	Barguna Sadar
	Galachipa	Galachipa	Galachipa	Galachipa	-	-	-	-

(d) Findings: Major Findings of Shalikhha Upazila

Physical Parameters

(a) Changes in Population over time: Naturally food requirement increased significantly.




Figure 1. Increase in population in Shalikha Upazila over time

(b) Changes in Settlements and Ponds: Area under settlement and ponds increased significantly.




Figure 2. Increase in settlement and ponds in Shalikha Upazila

(c) Temperature and rainfall: In the Shalikhha Upazila there are some changes in monthly maximum and minimum temperature observed considering long term average as the reference line. Maximum monthly temperature more than 30 degree Celsius was observed in the month of February to November. These changes might have effect on crop agriculture. There are some changes observed in annual rainfall and its distribution pattern. The long term average rainfall was 1,247 mm/year and maximum rainfall occurred in the month June to October. But in the recent years (2012-2016) it was observed that rainfall ranges from 1,305 to 1,867 mm/year. Total rainfall changes abruptly in every alternative years and maximum rainfall received in the month July to September. This might have effect on agriculture.

(d) Changes in Land Type

Land type	Previous Survey (1990)		Present Survey (2018)	
	Area (ha)	%	Area (ha)	%
Highland	4,217	18.4	2,934	12.8
Medium Highland	9,749	42.6	7,585	33.2
Medium Lowland	4,883	21.4	4,568	20.0
Lowland	467	2.1	391	1.7
Miscellaneous Land	3,548	15.5	7,386	32.3
Total	22,864	100.0	22,864	100.0



Field survey: Kazipur (left), Sirajdikhan (right)

(e) Changes in Land Use:

Land use types	Previous Survey (1990)		Present Survey (2018)	
	Area (ha)	%	Area (ha)	%
Perennial garden	256	1.1	38	0.2
Rabi vegetable-Kharif vegetable	85	0.4	239	1.1
Rabi crop-B.Aus/Jute-Fallow	2801	12.3	-	-
Rabi crop-Jute/ B.Aus /Til-T.Aman	1342	5.9	1179	5.2
Boro-Fallow-T.Aman	1124	4.9	6164	27.0
Rabi crop- Fallow-T.Aman	3058	13.4	361	1.6
Fallow-T.Aus/Jute- T.Aman	1999	8.7	180	0.8
Boro-B.Aman	1095	4.8	357	1.2
Rabi crop-Boro-T.Aman	-	-	2126	9.3
Boro-Deepwater T.Aman	999	4.4	-	-
Boro-Fallow-Fallow	2801	12.3	3302	14.4

Fallow-B.Aman	234	1.0	-	-
Rabi crop-B.Aman	1003	4.4	-	-
Other pattern	2604	11.4	639	2.8
Miscellaneous	3463	15.2	8279	36.2
Total	22864	100	22864	100

(e) Conclusion: Updating Upazila Land and Soil Resource Utilization Guide (Upazila Nirdeshika) through Semi-detailed soil survey is a vital program for understanding the changes in land use and soil fertility status of a specific area over time. This guide can be used as a modern tool for the transfer of the agricultural technology in national agricultural development.

3.2. Union Land, Soil and Fertilizer Recommendation Guide (Union Sahayika) Preparation

(a) Introduction: Union Land, Soil and Fertilizer Recommendation Guide (Union Sahayika) is being used as a tool for agricultural development activities at Union level. It is used to provide the existing information of Upazila Land and Soil Resource Utilization Guide (Upazila nirdeshika) to the farmers more intensely at Union level. From this guide, farmers can get land and irrigation information for their particular union and location specific fertilizer recommendations for the specific crops and varieties cultivated in that union. That is why it becomes imperative to convince the farmers for using the Sahayika for their economic benefit.

(b) Objectives

- To produce union level soil and land use database for local level agricultural development planning.
- To ensure the use of soil test based fertilizer recommendation at union level.
- Provide information regarding crop suitability and balanced fertilizer use at grass root level.

(c) Methodology

Base Materials: Existing Updated Upazila Soil and Landform Map (1:50,000).

Methods: Field survey is done at dry season. Base map (Upazila Soil and Landform Map) is divided into several grids and ground checking of each grid is done to verify changes. Land use data with other information is recorded in the profile card. Composite soil samples are collected and sent in the laboratory in the laboratory. After completing the survey map is prepared and with the result of soil analysis, Union Sahayika is prepared.

(d) Results and Discussion:

- 200 Union Sahayika has been prepared and published.
- Helped the farmers to use soil test based fertilizer doses in their fields.
- Now Sub Assistant. Agriculture Officers (SAAO) can easily help the farmers in advising farmers on fertilizer application.

(e) Conclusion: Union Sahayika is one of the important tools for using balanced fertilizer at union level. The Department of Agricultural Extension may play significant role in its extensive use.

3.3 Soil and Water Salinity Monitoring

Introduction

The coastal and offshore area of Bangladesh includes tidal, estuarine and meander river floodplains. The tidal floodplains occur in the south of the Ganges Floodplain and also on large parts of Chittagong coastal plains. The coastal and offshore area of Bangladesh includes tidal, estuarine and meander river floodplains. The tidal floodplains occur in the south of the Ganges Floodplain and also on large parts of Chittagong coastal plains. Salinity problem is now a

national concern. To explore the potentials of saline areas it is imperative to closely monitor the situation throughout the year. SRDI is regularly monitoring soil and water salinity in greater Khulna, Jessore, Barisal, Patuakhali, Noakhali and Chittagong districts.

Objectives

- To determine the soil and water salinity round the year in coastal area and to delineate area under different degrees of salinity.
- To determine the particular time frame in a year when surface water is suitable for irrigation.
- To record present land use and crop response to soil salinity.
- To provide necessary database which would help in developing appropriate reclamation technology and management practices.

Methodology

- Field data is being collected from strategically representative sites of the coastal area.
- Soil samples at variable depths (0-10cm, 10-30cm and 30-60cm), water samples from surface (river, canals, ponds, and waterbodies), underground water (Hand Tube Well, Shallow Tube Well, and Deep Tube Well) are collected twice in every month. Salinity of soils and water is determined by EC meter.

No. of Monitoring Sites

- Soil : 21 sites
- Water : 22 Sites
- DTW/HTW : 13 sites
- Total 560 samples collected and EC determined

Salinity condition of different rivers

1. Modhumoti river (Mollahat, Bagerhat) water salinity during 2012-2018

In 2018 water salinity was observed higher than the last 6 years, highest salinity being 4.0 dS/m in April. This river remains almost non-saline during June to December; water remains saline during January to May which is not suitable for irrigation (Figure 1).




Figure 3. Modhumoti river (Mollahat, Bagerhat) water salinity in during 2012-2018

2. Rupsa River (Khulna) water salinity during 2012-2018

In 2018 water salinity was observed lower than the last 2 years, highest salinity being 22.0 dS/m in April. This river remains almost non-saline during July to December, water remains saline during January to June and which is not suitable for irrigation (Figure 2).




Figure 4. Rupsha river (Khulna) water salinity in during 2012-2018

3. Water salinity of Bhadra river (Khornia, Dumuria, Khulna) during 2012-2018

In 2018 water salinity was observed higher than the last 6 years, highest salinity being 24.3 dS/m in April. This river remains almost non-saline during August to November, water remains saline during July to December and which is not suitable for irrigation (Figure 3).




Figure 5. Bhadra river (Khornia, Dumuria) water salinity in during 2012-2018

4. Water salinity of Betna river (Benerpota, Satkhira) during 2012-2018

In 2018 water salinity was observed lower than the last 2 years, highest salinity being 21.5 dS/m in April. This river remains almost non-saline during September to November; water remains saline during January to August which is not suitable for irrigation (Figure 4).




Figure 6. Betna river (Benerpota, Satkhira) water salinity in during 2012-2018

5. Water salinity of Kakshiali river (Kaligonj, Satkhira) during 2012-2018

In 2018 water salinity was observed lower than 2012 and 2015 but higher than other years, highest salinity being 28.5 dS/m in April. This river remains almost saline throughout the year and is not suitable for irrigation (Figure 5).




Figure 7. Kakshiali river (Kaligong, Satkhira) water salinity in during 2012-2018

6. Water salinity of Morichap river (Ashashuni, Satkhira) during 2012-2018

In 2018 water salinity was observed lower than last two years. The highest salinity being 27.2 dS/m in April. This river remains almost saline throughout the year and is not suitable for irrigation (Figure 5).




Figure 8. Morichap river (Ashashuni, Satkhira) water salinity in during 2012-2018

7. Water salinity of Shibsha river (Paikgachha, Khulna) during 2012-2018

In 2018 water salinity was observed lower than last two years. The highest salinity being 27.2 dS/m in April. This river remains almost saline throughout the year and is not suitable for irrigation (Figure 7).




Figure 9. Shibsha river (Paikgachha, Satkhira) water salinity in during 2012-2018

8. Water salinity of Kazibacha river (Batiaghata, Khulna) during 2012-2018

In 2018 water salinity was observed lower than last two years. The highest salinity being 25.0 dS/m in May. This river remains saline during July to November and remains saline during December to June which is not suitable for irrigation (Figure 8).




Figure 10. Kazibachha river (Batiaghata, Khulna) water salinity in during 2012-2018

9. Water salinity of Pasur river (Mongla, Bagerhat) during 2012-2018

In 2018 water salinity was observed lower than last two years. The highest salinity being 28.0 dS/m in May. This river remains saline during July to November and remains saline during December to June which is not suitable for irrigation (Figure 9).




Figure 11. Pasur river (Mongla, Bagerhat) water salinity in during 2012-2018

10. Water salinity of Daratana river (Bagerhat) during 2012-2018

In 2018 water salinity was observed lower than last six years. The highest salinity being 16.7 dS/m in May. This river remains saline during September to November and remains saline during December to August which is not suitable for irrigation (Figure 10).




Figure 12. Daratana river (Bagerhat) water salinity in during 2012-2018

11. Water salinity of Panguchhi river (Morelgonj, Bagerhat) during 2012-2018

In 2018 water salinity was observed higher than last six years. The highest salinity being 12.7 dS/m in March. This river remains saline during July to November and remains saline during December to June which is not suitable for irrigation (Figure 11).




Figure 13. Panguchhi river (Bagerhat) water salinity in during 2012-2018

3.4. Monitoring & Evaluation of Farmers' Service through Mobile Soil Testing Laboratories (MSTL)

(a) Introduction: Soil Test Based (STB) fertilizer use may be one of the ways to minimize the yield gap. The farmers in our country are not yet fully aware of the benefit of the STB fertilizer use in crop production. Moreover Static Laboratories are few in numbers and located district head quarters. The Government of Bangladesh has taken steps to provide analytical services at farmer's doorstep through MSTL. SRDI launched this program through two MSTL Program at block levels for specific location and yield goal basis fertilizer recommendation for crop based on STB since 1996. At present there are 10 MSTL for serving the farmer's of the country with soil testing facilities at Upazila level in Rabi and Kharif season every year.

(b) Objectives

- To create awareness among the farmers on benefit of using balanced fertilizer on the basis of STB and crop requirements.
- To serve the farmers with balanced fertilizer recommendation on the basis of STB & crop requirements by analyzing the soil samples at Upazila Sadar.
- To evaluate effect of soil test based fertilizer application on crop production.

(c) Methodology

About 180 farmers were selected for monitoring & evaluation under this program. The SRDI district offices monitor the activities of farmers and collect yield data with the help of DAE personnel. Yield was assessed by comparing the yield of the plots where fertilizers were applied following MSTL supplied Fertilizer Recommendation Cards (FRC) with adjacent plots having same crops and variety in the same season. The farmer's applied fertilizer dose is also recorded.

(d) Result and Discussion

In 2017-18, soil analytical service through MSTL was provided to 112 Upazila of the country and provided 5,600 farmers with Fertilizer Recommendation Cards (FRCs) (Table 2).

Table 2. Name of the Upazila and number of soil samples analyzed through MSTL under Farmer's Service Program during Rabi 2017 season and Kharif 2018 season.

Upazila	District	No. of Soil Samples Analyzed	Upazila	District	No. of Soil Samples Analyzed
Rabi 2018			Kharif 2018		
Manikganj Sadar	Manikganj	50	Daulatpur	Manikganj	46
Raipura	Narsingdi	50	Sreepur	Gazipur	43
Nawabganj	Dhaka	50	Palash	Narsingdi	46
Basail	Tangail	60	Tangail Sadar	Tangail	56
Delduar	Tangail	39	Bhuapur	Tangail	64
Goalanda	Rajbari	50	Char Bhadrashan	Faridpur	50
Saltha	Faridpur	43	Shibchar	Madaripur	48
Kalkini	Madaripur	43	Kalukhali	Rajbari	50
Gopalganj Sadar	Gopalganj	52	Kotalipara	Gopalganj	48
Mymensingh Sadar	Mymensingh	50	Purbadhala	Netrokona	33
Barhatta	Netrokona	52	Nandail	Mymensingh	43
Nikli	Kishoreganj	50	Katiadi	Kishoreganj	50
Jamalpur Sadar	Jamalpur	50	Melandaha	Jamalpur	50
Dewanganj	Jamalpur	50	Bakshiganj	Jamalpur	50
Madarganj	Jamalpur	50	Sherpur Sadar	Sherpur	50
Jhenaigati	Sherpur	50	Nalitabari	Sherpur	50
Laksam	Cumilla	50	Faridganj	Chandpur	61
Titas	Cumilla	50	Nangalkot	Cumilla	50
Sarail	Brahmanbaria	50	Ashuganj	Brahmanbaria	48
Noakhali Sadar	Noakhali	50	Feni Sadar	Feni	50

Upazila	District	No. of Soil Samples Analyzed	Upazila	District	No. of Soil Samples Analyzed
Rabi 2018			Kharif 2018		
Ramganj	Lakshmipur	50	Kabirhat	Noakhali	51
Sonagazi	Feni	47	Kamalnagar	Lakshmipur	50
Doublemooring	Chattogram	51	Fatikchhari	Chattogram	50
Kornofuli	Chattogram	51	Anwara	Chattogram	50
Boalkhali	Chattogram	50	Satkania	Chattogram	60
Pekua	Cox's Bazar	50	Kutubdia	Cox's Bazar	50
Rajasthali	Rangamati	51	Kawkhali	Rangamati	55
Bandarban Sadar	Bandarban	50	Matiranga	Khagrachhari	50
Mujibnagar	Meherpur	50	Mirpur	Kushtia	44
Alamdanga	Chuadanga	50	Gangni	Meherpur	47
Shailkupa	Jhenaidah	50	Harinakundu	Jhenaidah	48
Lohagara	Narail	50	Shalikha	Magura	50
Jhenaidah Sadar	Jhenaidah	51	Jessore Sadar	Jashore	50
Satkhira Sadar	Satkhira	59	Kaliganj	Satkhira	50
Dumuria	Khulna	43	Shyamnagar	Satkhira	50
Terokhada	Khulna	46	Koyra	Khulna	50
Debiganj	Panchagarh	81	Raiganj	Sirajganj	65
Boda	Panchagarh	64	Faridpur	Pabna	60
Kishorganj	Nilphamari	49	Dhamoirhat	Naogaon	39
Kaunia	Rangpur	52	Mohadevpur	Naogaon	81
Bera	Pabna	56	Mohonpur	Rajshahi	46
Bholahat	Chapainawabganj	50	Baliadangi	Thakurgaon	62
Gomostapur	Chapainawabganj	58	Birganj	Dinajpur	64
Nachole	Chapainawabganj	42	Sherpur	Bogura	51
Shibganj	Chapainawabganj	56	Joypurhat Sadar	Joypurhat	110
Panchbibi	Joypurhat	53	Gangachara	Rangpur	52
Belkuchi	Sirajganj	48	Bhurungamari	Kurigram	71
Bishwanath	Sylhet	50	Saghata	Gaibandha	34
Dakshin Sunamganj	Sunamganj	50	Lakhai	Habiganj	51
Barlekha	Moulvibazar	46	Moulvibazar Sadar	Moulvibazar	50
Chunarughat	Habiganj	68	Dakshin Surma	Sylhet	72
Nazirpur	Pirojpur	50	Derai	Sunamganj	42
Borhanuddin	Bhola	50	Barishal Sadar	Barishal	54
Galachipa	Patuakhali	50	Kathalia	Jhalakathi	50
Pathorghata	Barguna	50	Betagi	Barguna	51
			Dashmina	Patuakhali	50
Total		2,811	Total		2,900

In general farmers got higher yield of Transplanted Aus, Jute, Transplanted Aman, Brinjal, Cabbage, Potato, Water melon, Wheat, Mustard, Lentil, Indian spinach, Chilli, Corn, Tomato, Onion, Boro, Lemon if they use balanced fertilizer. In different locations, farmers' got 8 to 24% higher yield of BRRIdhan 29, 12 to 14% higher yield of BRRIdhan 28, 15 to 25% higher yield of mustard, 9 to 18% higher yield of wheat, 27 to 40% higher yield of potato by applying balanced fertilizer following FRC provided by MSTL in 2017-18.

Table 3. Comparative yield of FRC based fertilized crop and farmer's practice in Rabi 2017

Sl. No.	Name of Upazila	District	Crop & Variety	Average yield (t/ha)		
				Farmer's field	Demonstration plot	Yield increase
1	Manikganj Sadar	Manikganj	Boro BRRIdhan 29	6.25	7.5	16.6
	Keraniganj	Dhaka	Boro BRRIdhan 29	6.30	7.5	16
	Delduar	Tangail	Boro BRRIdhan 29	7.5	8.5	11.8
	Kalkini	Madaripur	Boro BRRIdhan 29	6.98	7.55	7.6
	Jamalpur Sadar	Jamalpur	Boro BRRIdhan 29	6.5	7.7	15.6
	Barhatta	Mymensingh	Boro BRRIdhan 29	5.7	7.0	18.6
	Gomostapur	Chapai-nawabganj	Rabi BARISharisha 14	1.2	1.5	25
	Gomostapur	Chapai-nawabganj	Rabi Bari Gom 26	3.45	4.05	17.4
	Mirzaganj	Patuakhali	Rabi Potato(Diamont)	28.4	21.4	32.7
	Mirzaganj	Patuakhali	Rabi Tomato (Hybrid)	32	40	25
	Bamna	Barguna	Rabi Potato (Diamond)	22	30	36

(e) Conclusion: Yields of Rabi crops increased from 7 to 32 percent due to application of balanced fertilizer following FRC. Motivation activities may help farmers' in using balanced fertilizers following Fertilizer Recommendation Cards provided by MSTL. This in turn will increase crop production in the country.

Chapter 4: Research Centers

4.1 Research activities of Soil Conservation and Watershed Management Center (SCWMC), Bandarban

PROGRAM 1. INTRODUCTION OF QUESUNGUAL SLASH AND MULCH AGRO-FORESTRY SYSTEM (QSMAS) FOR ENHANCING CROP YIELDS AND SOIL QUALITY IN CHITTAGONG HILL TRACTS

Abstract

The Chittagong Hill Tracts region is of great importance for growing various crops, which are different from the plains. Farmers practice traditional Jhum culture for their livelihood. They slash and burn the vegetation on hills and go for Jhum cultivation which contributes to soil and nutrient loss. It can also lead to land degradation if population pressure reduces the fallow periods needed for the recovery of natural resources. A Questionnaire based survey in six Upazilas of Bandarban revealed that the hill dwellers are lagging behind in education, farming knowledge, access to information on modern culture, inputs and loan facilities. As such they could seldom produce enough food for their consumption. The Quesungual Slash and Mulch Agro-forestry System (QSMAS) is a smallholder production system with a group of techniques for the sustainable management of vegetation, soil, and water resources in drought-prone hillsides is an alternative to traditional and widespread slash and burn agriculture. The present research work was undertaken to introduce an eco-friendly productive crop production system in sloping lands of CHT which will mitigate the process of land degradation due to Jhum culture as well as take care of food security of Hill people.

The experiment was conducted in moderate hill slope of Soil Conservation and Watershed Management Center, SRDI, Bandarban. Experimental site comprised of four plots-QSMAS model, modern Jhum with hedge row, traditional Jhum and control (secondary forest). From two years data it was observed that system productivity of QSMAS was much higher than that of Jhum with hedge row and traditional Jhum. Total soil loss and surface run off was much lower in QSMAS model than Jhum with hedge row and traditional Jhum. Soil organic matter and CEC was increased over initial status in QSMAS and secondary forest. Micro-watershed based agro-forestry system may be an alternate option to replace Jhum culture for livelihood security of Hill dwellers in Chittagong Hill Tracts.

INTRODUCTION

The Chittagong Hill Tracts comprising the three districts of Bandarban, Rangamati and Khagrachhari has an area of 13181sq km endowed with natural beauty and high economic potentiality. The tribal along with the Bengali people are living there for long maintaining their

distinct socio-cultural identities and harmony. The area is hilly with mild to very steep slopes (from 15% to over 70%) often breaking or ending in cliffs. More than 90 percent of the area is covered by hills with only 129,000 hectares of cropped land. About 87 per cent of the land is covered with forest (totaling 11,475 sq.km) mostly owned by the government (Dasgupta and Ahmed, 1998). Presently, it is increasingly becoming denuded due to unplanned management of hills and agricultural practices at steep slope without any conservation measure. There are hills with altitudes of more than 3000 feet (Brammer, 1986) having steep and long slope. The total annual precipitation is also high (2000-3550mm). Continuous depletion of soil fertility is the major constraint to sustainable crop production in the hilly areas of Bangladesh. According to Banglapedia (2009) about 20,000 hectares of land are being brought under Jhum cultivation every year.

Jhum cultivation, sloppiness, heavy rainfall and improper management of soil enhanced nutrient depletion through erosion. Accelerated soil erosion is the greatest hazard for the long term maintenance of soil fertility. Gafur *et al.* (2003) carried out a research to find out runoff and losses of soil and nutrients from small watersheds under shifting cultivation in the CHT. Borggaard *et al.* (2003) carried out a study to analyze the sustainability appraisal of shifting cultivation in CHT. Dewan (2008) conducted a survey work to analyze the socio-economic status of Jhum cultivators. The Chittagong Hill Tracts region is of great importance for growing various crops, which are different from the plains. But unfortunately few eco-friendly sustainable agriculture practices for CHT have so far been developed.

Slash and burn practices, also known as *shifting cultivation*, *swidden agriculture*, or simply *jhum chash*, is an ancient form of agriculture practiced by 200 to 500 million people around the world currently. The people in the uplands of eastern Bangladesh have been practicing shifting cultivation from time immemorial and it is closely related with their socio-cultural identity (Miah and Islam, 2007). However, in the past, they practiced shifting cultivation in the same area with a fallow period of 15–20 years, which ensured the long-term sustainability of soil fertility, and ensured forest regrowth. With the rapid growth in population, the fallow period has been dramatically reduced to 3–4 years, allowing very little time for soil or vegetative regeneration (Riessen, 2000). The decrease in fallow period has led to the deterioration of faunal and microbial organisms, top soil loss, and erosion during periods of heavy rainfall (Gafur, 2001).

The two key components of slash and burn agriculture are the use of fire to prepare fields for cultivation and the subsequent abandonment of those fields as productivity declines. The inevitable decline in productivity is a result of the depletion of soil nutrients and also a result of the invasion of weed and pest species (Cornell, 2011). Slash and burn contributes to global warming by acting as a major source of greenhouse gas emissions, and by depleting reserves of carbon both above and below-ground. It can also lead to land degradation if population pressure reduces the fallow periods needed for the recovery of natural resources. With the increasing population pressure several alternatives to shifting cultivation have been suggested (FAO, 1984)

which include: (1) tree crop plantation, (2) agro-forestry, (3) planted fallow system (tree and shrub fallows plus arable crop sequence), (4) livestock production, and (5) special commercial horticulture.

Eco-efficient agriculture uses resources more efficiently to achieve sustainable increases in productivity, reduces the degradation of natural resources, and creates opportunities for boosting incomes and employment in rural areas. The Quesungual Slash and Mulch Agro-forestry System (QSMAS) is one example of eco-efficient crop production for tropical sub-humid regions. It has reduced erosion and improved crop yields and quality of life for over 6,000 local families while allowing regeneration of about 60,000 hectares of secondary forest (*New Agriculturalist* 2009).

QSMAS is a smallholder production system with a group of techniques for the sustainable management of vegetation, soil, and water resources in drought-prone hillsides. The system was developed in the early 1990s in close collaboration with farmers and technicians from FAO and other institutions, as an alternative to traditional and widespread slash and burn agriculture. It has had an extraordinary impact on the livelihoods of farmers growing maize, beans, and sorghum in Central America, and has great potential to be used in other regions.




Figure 27. Site similarity analysis: bivariate map showing potential areas for targeting QSMAS across the developing countries in the tropics (performed by combining Bayesian and frequency probability statistical analyses)

Past research reports indicate that little work has been undertaken so far on replacing the traditional Jhum system with modern techniques to reduce soil erosion, biodiversity loss, deforestation, factors that contribute to environmental degradation and impacts on environment due to shifting (Jhum) cultivation practice.

Keeping the above views in mind the present research work was undertaken to introduce a eco-friendly productive crop production system in sloping lands of CHT which will mitigate the process of land degradation due to Jhum culture as well as take care of food security of Hill people.

Goal: Introduce an eco-efficient crop production system in sloping lands of CHT

Objective (s):

- i) Establish Quesungual Slash and Mulch Agro-forestry System (QSMAS) in CHT.

- ii) To improve both the productivity and economic returns of land currently engaged in slash and burn agriculture.
- iii) To create awareness about soil conservation and watershed management among hill dwellers.

MATERIALS AND METHODS

Eco-efficient agriculture uses resources more efficiently to achieve sustainable increases in productivity, reduces the degradation of natural resources, and creates opportunities for boosting incomes and employment in rural areas.

To validate the principles of Quesungual agro-forestry system in Soil Conservation and Watershed Management Centre, SRDI, Bandarban watershed four land use systems were established: traditional Jhum (slash-and-burn), Jhum with modern management, Quesungual slash and mulch agro-forestry systems (QSMAS), and demarcated areas of secondary forest as a control. Crops like rice, maize; millet, cotton, sesame and common beans, marfa, yard long bean, sweet gourd, ginger and turmeric were accommodated in a traditional system, application of slashed vegetation/crop residues as mulch and QSMAS, to measure and compare differences among production systems. Soil sampling for initial fertility assessment and determine change in fertility status after each cropping season for three years.

Soil sampling consisted of digging test pit of 50 cm depth and sampling of soil at 0-13, 13-43, 43-63 cm depths just before sowing every year. Composite soil samples will be collected from each plot for fertility determination. Chemical characterization included determination of pH, organic matter (OM), N, P, K, S, Zn, B, Ca, Mg, Mn, Fe, Cu. In the field, productivity of rice, maize; cotton, sesame and common beans, marfa, sweet gourd, ginger and turmeric were evaluated in three cropping seasons, from 2014 to 2017.

Statistical analyses of soil fertility and crop yields data were done to determine the change in soil fertility and crop productivity over the years.

Layout of experimental plots

Experimental site: SCWMC, Meghla, Bandarban




Figure 28. Layout of experimental site (moderate slope; Plot size (5mX20m) =100m²)

RESULTS AND DISCUSSION

SOIL FERTILITY STATUS

Initial fertility status was compared with fertility status of each plot after crop harvest. Soils are mostly highly acidic to slightly acidic in nature. Initial Organic matter status was low to medium while it was high to medium after crop harvest. Initial Nitrogen status was very low to low while it was low to medium after crop harvest. Phosphorus status was very low. Initial Potassium status is medium to optimum while it was medium to very high after crop harvest. Sulfur status was reduced from medium to low to low to very low. Zinc status was also reduced from initial status i.e. low to medium. Boron status reduced from very high to low to medium. Calcium, Magnesium, Copper, Iron and Manganese status is remained almost unchanged (Table 1). Physical analysis was done to determine the soil texture (Table 2). CEC, which indicates soil fertility, seems to be improved over time under QSMAS and secondary forest system. Highest CEC increase was observed in QSMAS plot (Figure 15). Whereas, CEC was reduced in Jhum with hedge and Traditional Jhum system.

Table 4. Initial soil fertility status and fertility status after crop harvest

Plot No./ Year	Depth of soil sample	pH	OM (%)	N (%)	P	K	S	Zn	B	Ca	Mg	Cu	Fe	Mn
					meq/100g soil	µg/g soil			meq/100g soil	µg/g soil				
1/2015	0-13	4.9	1.82	0.10	2.85	0.33	15.0	1.10	0.93	2.53	1.40	1.06	75.93	15.18
			M	L	VL	O	M	M	VH	L	O	VH	VH	VH
1/2016	0-13	4.5	4.13	0.24	5.32	0.52	6.05	1.01	0.18	5.82	2.46	0.72	71.80	11.22
			H	M	VL	VH	VL	M	L	O	VH	H	VH	VH
2/2015	0-13	5.7	1.62	0.09	1.21	0.35	8.17	0.64	0.86	1.77	1.16	0.81	76.28	12.67
			L	VL	VL	O	L	L	VH	L	M	VH	VH	VH
2/2016	0-13	4.5	2.88	0.17	3.88	0.26	6.08	0.09	0.22	2.20	1.08	0.42	11.72	2.34
			M	L	VL	M	VL	VL	L	L	M	M	O	H
3/2015	0-13	4.9	1.32	0.07	1.38	0.32	9.17	0.88	0.92	3.21	1.37	0.99	86.34	10.20
			L	VL	VL	O	L	L	VH	M	O	VH	VH	VH
3/2016	0-13	4.5	3.12	0.18	6.86	0.42	7.35	0.96	0.16	2.50	1.17	0.44	43.8	14.3
			M	L	VL	H	VL	M	L	L	O	M	VH	VH
4/2015	0-13	4.9	1.10	0.06	2.19	0.26	5.20	0.78	0.79	2.18	1.16	0.88	81.15	7.80
			L	VL	VL	M	VL	L	VH	L	M	VH	VH	VH
4/2016	0-13	4.5	3.24	0.19	6.80	0.46	8.42	1.02	0.21	1.25	1.04	0.14	42.2	2.80
			M	M	VL	VH	L	M	M	VL	M	VL	VH	O

Note: VL=very low; L=low; M= medium; O=optimum; VH=very high

Table 5. Mean, standard error, correlation coefficient and significance of soil fertility indicators over time

Soil nutrients	Mean ± SE		Correlation	significance
	2015	2016		
pH	5.10±0.20	4.50±0.00	0.00	0.00
OM	1.47±0.16	3.34±0.27	0.53	0.47
N	0.08±0.01	0.20±0.02	0.53	0.47
P	1.91±0.38	5.72±0.71	0.19	0.81
K	0.32±0.02	0.42±0.06	-0.49	0.51
S	9.39±2.05	6.98±0.57	-0.73	0.27
Zn	0.85±0.10	0.77±0.23	0.72	0.28
Ca	2.42±0.31	2.94±1.00	0.24	0.77
Mg	1.27±0.07	1.44±0.34	0.71	0.29
Cu	0.94±0.06	0.43±0.12	0.68	0.32
Fe	79.93±2.45	42.38±12.27	0.01	0.99
Mn	11.46±1.59	7.67±3.01	0.28	0.72




Figure 29. Comparative CEC data of experimental plots over time

Table 6. Soil texture analysis data

Plot No.	Sampling depth (cm)	Soil texture	Sand (%)	Silt (%)	Clay (%)
1	0-13	Sil loam	17	59	24
2	0-13	Silt loam	19	59	22
3	0-13	Silt loam	20	58	22
4	0-13	Silt loam	18	58	24
Layer wise sample					
	0-13	Silt loam	13	61	26
	13-43	Clay loam	24	44	32
	43-63+	Clay loam	21	40	39

Crop yield and system productivity

Each plot had the same crop combination except control (secondary forest). Rice (local), maize (local), sesame, millet, sweet gourd, chilly, marfa, yardlong bean, country bean, cotton, ginger, turmeric were planted in mixture. But in QSMAS model the crops were arranged in sub plots within the main plot. Grafted fruit trees-mango, carambola and seedlings of papaya were planted in the plot. After harvestings crop yield data were recorded and analyzed. It was observed that rice yield was higher in traditional Jhum practice than other practices. But system productivity of QSMAS model plot was much higher than other plots (Table 4, 5 & 6). Comparative yield and economic data of component crops are shown in Tables 7 and 8.

Table 7. Yield (kg/100 sqm) and return (BDT) of crops harvested from experimental plots (2015)

Sl.No	Crops	Yield (kg/100 sqm)			Price (BDT/Kg)	Return in BDT			Remarks
		Traditional Jhum	Jhum with hedge row	QSMAS model		Traditional Jhum	Jhum with hedge row	QSMAS model	
1	Rice (local)	15.00	10.00	6.00	15.00	225.00	150.00	90.00	QSMAS model out yielded all the other plots
2	Maize (local)	3.00	5.00	6.00	50.00	150.00	250.00	300.00	
3	Sesame	1.00	1.50	1.50	60.00	60.00	90.00	90.00	
4	Millet	1.00	0.50	0.60	80.00	80.00	40.00	48.00	
5	Sweet gourd	4.00	5.00	6.00	35.00	140.00	175.00	210.00	
6	Chilly	0.40	0.50	0.50	120.00	48.00	60.00	60.00	
7	Marpha	3.00	4.00	4.00	40.00	120.00	160.00	160.00	
8	Yardlongbean	4.00	5.00	6.00	40.00	160.00	200.00	240.00	
9	Countrybean	-	-	6.00	60.00	-	-	360.00	
10	Cotton	1.50	2.00	3.00	200.00	300.00	400.00	600.00	
11	Ginger	3.00	5.00	7.00	60.00	180.00	300.00	420.00	
12	Turmeric	12.00	15.00	17.00	20.00	240.00	300.00	340.00	
13	Mango (4)							-	No fruiting was observed
14	Papaya (5)							-	
15	Carambola (3)							-	
Total						1703.00	2125.00	2918.00	

Table 8. Yield (kg/100 sqm) and return (BDT) of crops harvested from experimental plots (2016)

Sl.No.	Yield (kg/100 sq m)			Price (BDT/Kg)	Return in BDT			Remarks	
	Crops	Traditional Jhum	Jhum with hedge row		QSMAS model	Traditional Jhum	Jhum with hedge row		QSMAS model
1	Rice (local)	14.00	13.00	8.00	15.00	210.00	195.00	120.00	QSMAS model out yielded all the other plots
2	Maize (local)	3.00	5.00	4.00	50.00	150.00	250.00	250.00	
3	Sesame	0.80	1.00	0.90	60.00	48.00	60.00	54.00	
4	Millet	0.90	0.80	0.50	80.00	72.00	64.00	40.00	
5	Sweet gourd	3.50	4.00	5.00	35.00	122.00	140.00	175.00	
6	Chilly	1.20	1.00	0.70	80.00	96.00	80.00	56.00	
7	Marpha	2.50	4.00	3.00	40.00	100.00	160.00	120.00	
8	Yardlongbean	5.00	9.00	6.00	35.00	175.00	315.00	210.00	
9	Countrybean	-	-	4.00	40.00	-	-	160.00	
10	Cotton	1.00	1.50	1.20	200.00	200.00	300.00	240.00	
11	Ginger	3.00	6.00	4.00	50.00	150.00	300.00	200.00	
12	Turmeric	12.00	17.00	14.00	10.00	120.00	170.00	140.00	
13	Mango (4)	-	4 nos.	2.00	80.00	-	-	160.00	
14	Papaya (5)	-	5 nos.	40.00	15.00	-	-	600.00	
15	Carambola (3)	-	3 nos.	3.00	10.00	-	-	30.00	
Total						1443.00	2034.00	2505.00	

Table 9. Yield (kg/100 sqm) and return (BDT) of crops harvested from experimental plots (2017)

Sl.No.	Yield (kg/100 sq m)			Price (BDT/Kg)	Return in BDT			Remarks	
	Crops	Traditional Jhum	Jhum with hedge row		QSMAS model	Traditional Jhum	Jhum with hedge row		QSMAS model
1	Rice (local)	9.00	10.00	7.00	22.00	198.00	220.00	154.00	QSMAS model out yielded all the other plots
2	Maize (local)	2.00	4.00	5.00	50.00	100.00	200.00	250.00	
3	Sesame	0.70	0.80	1.00	60.00	42.00	48.00	60.00	
4	Millet	0.60	0.80	0.90	40.00	24.00	32.00	36.00	
5	Sweet gourd	3.00	5.00	6.00	35.00	105.00	175.00	210.00	
6	Chilly	0.50	0.70	0.90	80.00	40.00	56.00	72.00	
7	Marpha	2.00	3.00	4.50	40.00	80.00	120.00	180.00	
8	Yardlongbean	3.00	5.00	6.00	35.00	105.00	175.00	210.00	
9	Countrybean	-	-	5.00	40.00	-	-	200.00	
10	Cotton	0.60	0.80	1.00	200.00	120.00	160.00	200.00	
11	Ginger	2.50	3.50	4.00	50.00	125.00	175.00	200.00	
12	Turmeric	9.00	14.00	12.00	10.00	90.00	140.00	120.00	
13	Mango (4)	-	4.00	7.00	80.00	320.00	320.00	560.00	
14	Papaya (5)	-	20.00	26.00	15.00	300.00	300.00	390.00	
15	Carambola (3)	-	5.00	7.00	10.00	50.00	50.00	30.00	
Total						1699.00	2171.00	2912.00	

Table 10. Comparative yield of component crops over time

Crops	Mean \pm SE			Correlation	Sig.
	2015	2016	2017		
Rice	10.33 \pm 2.60	11.67 \pm 1.86	8.67 \pm 0.88	0.61	0.22
Maize	4.67 \pm 0.88	4.00 \pm 0.58	3.67 \pm 0.88	1.00	0.85
Sesame	1.33 \pm 0.17	0.90 \pm 0.17	0.83 \pm 0.09	0.76	0.30
Millet	0.70 \pm 0.16	0.73 \pm 0.12	0.77 \pm 0.09	-0.87	0.43
Sweet gourd	5.00 \pm 0.58	4.17 \pm 0.44	4.67 \pm 0.88	0.98	0.29
Chilli	0.47 \pm 0.03	0.97 \pm 0.15	0.70 \pm 0.12	0.87	0.41
Marpha	3.67 \pm 0.33	3.17 \pm 0.44	3.17 \pm 0.73	0.80	0.42
Yardlong bean	5.00 \pm 0.58	6.67 \pm 1.20	4.67 \pm 0.88	0.98	0.65
Cotton	2.17 \pm 0.44	1.90 \pm 0.56	0.80 \pm 0.12	0.93	0.90
Ginger	5.00 \pm 1.15	3.67 \pm 1.45	3.33 \pm 0.44	0.89	0.77
Turmeric	14.67 \pm 1.45	14.33 \pm 1.45	11.67 \pm 1.45	0.68	0.87

Table 11. Comparative return (BDT) from component crops over time

Crops	Mean ± SE			Correlation	Sig.
	2015	2016	2017		
Rice	155.00± 39.05	175.00± 27.84	190.67±19.40	0.61	0.22
Maize	233.33± 44.10	216.67± 33.33	183.33±44.10	1.00	0.85
Sesame	80.00± 10.00	54.00± 3.46	50.00±5.29	0.76	0.30
Millet	56.00± 12.22	58.67± 9.61	30.67±3.53	-0.87	0.43
Sweet gourd	175.00± 20.21	145.67± 15.56	163.33±30.87	0.98	0.39
Chilli	56.00± 4.00	77.33± 11.62	56.00±9.24	0.87	0.41
Marpha	146.67± 13.33	126.67± 17.64	126.67±29.06	0.80	0.42
Yardlong bean	200.00± 23.09	233.33± 42.06	163.33±30.87	0.98	0.67
Cotton	433.33± 88.19	246.67± 29.06	160.00±23.09	0.89	0.80
Ginger	300.00± 69.28	216.67± 44.10	166.67±22.05	0.93	0.67
Turmeric	293.33± 29.06	143.33± 14.53	116.67±14.53	0.68	0.87

Soil loss from experimental plots

As Bandarban is a high rainfall area if the soil surface is exposed due to deforestation it becomes vulnerable to water erosion. Soil loss from hills depends on surface cover, rainfall intensity, nature of slope and aspects of slope. Bandarban experienced a significant amount of rain every year though its distribution uneven over months. Rainfall intensity is higher in the months of May to August (Figure 16). Multi-slot divisor was established at the bottom of each plot. Total surface run-off and total soil loss was calculated per shower and cumulative figure was made by adding each observation. Last three years it was observed that highest total soil loss (39.17t ha⁻¹ y⁻¹-2015) occurred in traditional Jhum plot followed by Jhum with hedge and mulch and QSMAS model.




Figure 30. Monthly distribution of rainfall in experimental area over time


The lowest total soil loss was observed in control plot (secondary forest). In 2016 similar trends were observed. The finding is in conformity with that of CIAT (2010). Gafur *et al.* (2003) conducted a research to find out the runoff and losses soil and nutrients from small watersheds under shifting cultivation in the Chittagong Hill Tracts. In similar studies, Shoaib *et al.*(1998) recorded total soil loss to be 40-45t ha⁻¹y⁻¹ in traditional Jhum culture highest being observed in steep slope and the lowest in gentle slope. There is an evidence that the use of contour hedgerows on steep slopes (40-50%) can reduce erosion by 55-80% and run off by 30-70%

compared to shifting cultivation (Khisa, 2001). It was observed that QSMAS protects soil by markedly reducing soil erosion (Figure 17) in comparison to Jhum plots. This result is in conformity with the findings of CIAT (2010).

Table 12. Total soil loss from experimental plots ($t\ ha^{-1}\ y^{-1}$) in 2015, 2016 and 2017

Particulars		Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Total ($t\ ha^{-1}\ y^{-1}$)
Control	2015	-	-	-	-	0.80	3.32	3.12	3.14	1.09	0.67	-	-	12.14
	2016	-	-	-	-	-	2.28	1.53	0.83	0.52	0.63	-	-	5.79
	2017	-	-	-	-	0.45	1.12	1.47	0.82	0.51	0.72	0.5	-	5.59
QASMAS	2015	-	-	-	-	1.68	6.18	4.52	4.49	1.52	1.65	-	-	20.04
	2016	-	-	-	-	-	4.55	2.57	1.63	0.96	0.72	-	-	10.43
	2017	-	-	-	-	1.11	1.98	2.67	1.42	1.04	1.10	0.79	-	10.11
Jhum with hedge row	2015	-	-	-	-	2.15	7.84	5.58	5.67	1.96	1.90	-	-	25.10
	2016	-	-	-	-	-	7.01	4.34	1.89	1.06	0.86	-	-	15.16
	2017	-	-	-	-	1.64	3.14	3.54	1.92	1.68	1.84	1.30	-	15.06
Traditional Jhum	2015	-	-	-	-	2.68	10.52	9.18	9.49	4.07	3.23	-	-	39.17
	2016	-	-	-	-	-	9.4	8.4	2.77	1.25	1.36	-	-	23.18
	2017	-	-	-	-	2.55	4.40	5.80	2.82	2.46	2.86	2.41	-	23.30

Total soil loss from different plots(ton/ha)



It was observed that highest total soil loss occurred in traditional Jhum plot followed by Modern Jhum (with mulch) and QSMAS model. The lowest total soil loss was observed in control plot (secondary forest). The finding is in conformity with that of CIAT (November 2010).

Figure 31. Reduction of soil loss over time under different land use systems




Figure 32. QSMAS protects soil by markedly reducing soil erosion (87%, 84% and 67% less after 1, 2 and 3 years, respectively) compared to slash and burn

Generally, $11.2 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ is considered as permissible erosion limit based on the assumption that this rate of erosion equals soil formation. However, the actual erosion occurring in different sites sometimes goes far beyond this assumption. Thus, considering a single value approach as critical limit for soil erosion would be misleading (Lakaria *et al.* 2010)

Runoff and sediment load

The total runoff per hectare during 2015, 2016 and 2017 cropping season was highly variable between experimental plots, although there was no difference in terms of the rainfall received during the same period. The distribution of runoff during the years is shown in Table 11 as monthly values. The distribution of runoff follows the rainfall amount and intensity pattern with the maximum monthly runoff occurring in June, irrespective of land use. On average, the highest runoff volume was from traditional Jhum. The runoff from the watersheds and the sub-watersheds seems to have been influenced by factors such as topographic characteristics, land use and management practices implemented (Hartantoa *et al.*, 2003; Gary and Carmen, 2007).

Table 13. Total surface run off (%) from experimental plots in 2015, 2016 and 2017

Particulars		Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec
Control	2015	-	-	-	-	4.40	35.61	65.30	41.74	47.40	17.73	-	-
	2016	-	-	-	-	-	35.92	29.65	30.17	29.40	50.43	-	-
	2017	-	-	-	-	7.07	36.69	17.69	18.17	14.27	11.73	15.24	-
QASMAS	2015	-	-	-	-	4.87	46.05	67.41	51.63	57.71	21.80	-	-
	2016	-	-	-	-	-	41.17	47.90	39.53	30.84	35.63	-	-
	2017	-	-	-	-	8.42	41.14	20.31	21.29	16.18	13.82	17.61	-
Jhum with hedge row	2015	-	-	-	-	5.18	50.31	69.22	60.62	66.73	22.82	-	-
	2016	-	-	-	-	-	44.73	55.74	47.30	32.28	40.57	-	-
	2017	-	-	-	-	9.43	43.57	22.40	22.86	20.02	16.60	19.97	-
Traditional Jhum	2015	-	-	-	-	5.87	52.19	71.03	72.90	75.76	23.84	-	-
	2016	-	-	-	-	-	49.20	64.54	51.65	33.72	45.50	-	-
	2017	-	-	-	-	10.45	48.02	25.80	24.42	25.77	17.99	22.34	-

Nutrient loss through surface runoff and erosion

The impact of soil erosion on the productive potential of agricultural lands is well known (Pathak *et al.*, 2004), but the magnitude depends on local circumstances. In the study areas, the organic matter depletion was also observed irrespective of land use. The loss of the essential plant nutrients (N, K, S, Zn, B, Ca, Mg and Mn) in association with the suspended sediments and runoff during the measurement period was remarkable (Figures 19 through 27). The selective erosion of plant nutrients in runoff is a well known phenomena (Sharpley, 1985), and the sediment lost from the experimental plots on the micro-watershed was clearly enriched in all elements except P, relative to the topsoil of the watershed. The highest loss was displayed by Mn, Zn and S possibly resulting from reductive dissolution of oxides caused by sudden saturation of the soils in the earlier heavy rainfalls of the season. The results are in partial conformity with Gafur *et al.* (2003). This suggests that soil conservation control efforts should be prioritized in areas with high soil and nutrient loss potential so that their productivity is maintained.

Table 14. Nutrient loss (tha^{-1}) from plots under different land use (2016)

Particulars	N	P	K	S	Zn	B	Ca	Mg	Cu	Mn
Secondary forest	5.40	0.04	0.65	0.88	0.18	0.01	5.1	0.25	0.024	0.093
QSMAS	4.00	0.04	0.44	0.54	0.16	0.02	2.0	0.21	0.031	0.178
Jhum with hedge	4.80	0.60	0.26	0.94	0.17	0.01	2.1	0.19	0.031	0.154
Traditional Jhum	4.80	0.04	0.26	0.56	0.17	0.01	1.7	0.26	0.028	0.193

It was observed that highest nitrogen loss i.e. 5.4 tha^{-1} occurred from secondary forest plot and the lowest (4.0 tha^{-1}) from QSMAS plot along with other nutrient elements. In case of Jhum with hedge and Traditional Jhum plot nitrogen loss was 4.8 tha^{-1} along with other nutrient elements. Gafur (2001) found that in each year, the eroded soil from all the Jhum fields in CHT carries out about 4,309 tons of nitrogen along with other nutrients and about 14,071 tons of commercial fertilizers would be required to replace nutrients in eroded soil that would cost approximately US \$1.8 million annually.



Photo: Slash and mulch agro-forestry at SCWMC, Bandarban

Figure 33. Dynamics of OM status due to soil erosion




Figure 34. Dynamics of Total N due to soil erosion




Figure 35. Dynamics of P status due to soil erosion




Figure 36. Dynamics of K status due to soil erosion




Figure 37. Dynamics of S status due to soil erosion




Figure 38. Dynamics of Zn status due to soil erosion




Figure 39. Dynamics of B status due to soil erosion




Figure 40. Dynamics of Ca status due to soil erosion




Figure 41. Dynamics of Mg status due to soil erosion





Figure 42. Dynamics of Mn status due to soil erosion



CONCLUSION

A Questionnaire based survey in six Upazilas of Bandarban revealed that the hill dwellers are lagging behind in education, farming knowledge, access to information on modern culture, inputs and loan facilities. As such they could seldom produce enough food for their consumption. Fertility status was observed to be improved due to improved land use and management system.

System productivity of QSMAS plot was much higher than that of other plots. It was observed that highest total soil loss occurred in traditional Jhum plot followed by Jhum with hedge row and QSMAS model. The lowest total soil loss was observed in control plot (secondary forest).

Therefore, effective measures should be taken to discourage slash and burn shifting cultivation and upscale farmers knowledge base and information for awareness building along with other supports to enable them to adopt agro-forestry as a sustainable land-use system.

Agro-forestry is considered as one of the major strategies for sustainable forest management as well as poverty reduction in Bangladesh, where there is obvious priority for food crop production. Farmers need technical know-how, capital investment, marketing facilities and institutional supports (mainly title to their land) to move from Jhum to sedentary farming practices such as agro-forestry. Micro-watershed based agro-forestry system may be an alternate option for Jhum culture for livelihood security of Hill dwellers in Chittagong Hill Tracts. Government should design necessary program to address the needs of upland farmers, e.g., substantial initial investment, to support them to move from shifting cultivation to agro-forestry systems.

RECOMMENDATION

Effective measures should be taken to discourage Slash and burn cultivation and upscale farmers knowledge base and information for awareness building along with other supports to enable them adopt slash and mulch agro forestry as a sustainable land use system.

REFERENCES

- Asian Development Bank (ADB) (2001). Chittagong Hill Tracts region development plan. Final report, ADB TA no. 3328, Rangamati, Bangladesh.
- Bangladesh-Household Income and Expenditure Survey.2005. (BGD) HIES v01 M. Bangladesh Bureau of Statistics, Sept 29, 2011.
- Banglapedia(2009).Population-
Banglapedia.<http://en.banglapedia.org/index.php?title=Population>
- Borggaard, Ole K., Gafur, A. and Peterson, L. (2003) Sustainability appraisal of shifting cultivation in the Chittagong Hill Tracts of Bangladesh. *Ambio*. 32 (2), 118-123.<https://doi.org/10.1579/0044-7447-32.2.118>
- Brammer H (1986) Reconnaissance soil and land use survey: Chittagong Hill Tracts (1964–1965). Soil Resources Development Institute, Bangladesh
- Chakma, A. S. and Nahar, B. S. (2012). Jhum cultivation influences the degradation of hilly environment. *Journal of Environmental Science & Natural Resources*, 5(2), 339-344. www.banglajol.info/index.php/JESNR/article/view/14840
- Dasgupta, S. and Ahmed, F. U. (1998). Natural resource management by tribal community: a case study of Bangladesh. pp. 52-55.
- de Rouw, A. 1995. The fallow period as a weed-break in shifting cultivation (tropical wet forests). *Agric. Ecosyst. Environ.* 54: 31–43. [https://doi.org/10.1016/0167-8809\(95\)00590-O](https://doi.org/10.1016/0167-8809(95)00590-O)
- Dewan (2008). Conducted a survey work to analyze the socio-economic status of 70 jhum cultivators. This was quoted by Chakma,AS and Nahar,BS 2012 in their publication but I failed to locate original paper, so please drop this statement along with reference.
- Dingkuhn, M., Johnson, D. E., Sow, A. and Audebert, A. Y. (1999). Relationships between upland rice canopy characteristics and weed competitiveness. *Field Crops Res.* 61, 79-95.
- Gafur, A., Borggaard, O.K., Jensen, J.R., Peterson, L. 2003. Run off and losses of soil and nutrients from watershed under shifting cultivation (Jhum) in the Chittagong Hill Tracts of Bangladesh. *J. Hydrol.*, 279:293-309. [https://doi.org/10.1016/S0022-1694\(03\)00263-4](https://doi.org/10.1016/S0022-1694(03)00263-4)
- Gafur A (2001) Effects of shifting cultivation on soil properties, erosion, nutrient depletion, and hydrological responses in small watershed of the Chittagong Hill Tracts of Bangladesh. PhD Dissertation, The Royal Veterinary and Agricultural University, Copenhagen Household Income and Expenditure Survey Analysis Report. 2005. Bangladesh Bureau of Statistics, Dhaka, Bangladesh.

- Gary, W. C. and Carmen, V.: Impacts of land use changes on runoff generation in the east branch of the brandywine creek watershed using a GIS-based hydrologic model, *Middle States Geographer*, 40, 142–149, 2007.
- Hartantoa, H., Prabhub, R., Widayat, A., and Asdakd, C.: Factors affecting runoff and soil 30 erosion: plot-level soil loss monitoring for assessing sustainability of forest management, *Forest Ecol. Manag.*, 180, 361–374, 2003.
- Jamaluddin, M., Hassan, M. K. & Miah, M. M. 2010. Identifying livelihood patterns of ethnic minorities and their coping strategies different vulnerabilities situation in Chittagong hill tracts region, Bangladesh. Final Report. National Food Policy Capacity Strengthening Programme Food Planning and Monitoring Unit (FPMU), Ministry of Food and Disaster Management. Government of Bangladesh. p. 53.
- Lakaria, B. L., Mandal, D., Biswas, H. 2010. Permissible soil erosion limits under different landscapes of Chhattisgarh. *Indian Journal of Soil Conservation* 2010 Vol. 38 No. 3 pp. 148-154
- Miah MAM, Islam SMF (2007) Shifting cultivation and its alternatives in Bangladesh: productivity, risk and discount rates. SANDEE working paper no. 24-07, Nepal
- Pathak, P., Wani, S. P., Singh, P., and Sudi, R.: Sediment flow behavior from small agricultural watersheds, *Agr. Water Manage.*, 67, 105–117, 2004.581
- Rasul, G. and Thapa, G. B. (2002). State policies, praxies and land use in chittagong hill tracts of Bangladesh, *Regional and Rural Development planning*, School of Environment, Resource and Development, Asian Institute of Technology, Thailand, Bangkok.
- Riessen AV (2000) Chittagong Hill Tracts region development plan. Interim report no. 7. *Sociology and Indigenous People*, Asian Development Bank, Bangladesh
- Shoaib, J.U., Mostafa, G., Rahman, M. 1998. A Case Study on Soil Erosion Hazard in Hilly Regions of Bangladesh, Annual Report, SRDI. Dhaka.
- Tripura, P. and Harun, A. (2003). Parbotto Chattagrame Jum chash.
- Ullah, M. M., Malek, M. A., Karim, M. M. & Ali, M. S. (2012). A report on jhum research in CHT. Hill Agriculture Research Station, Bangladesh Agricultural Research Institute, Khagrachhari Hill District. p. 31.
- Van, K. K. (2003). On-farm agronomic diagnosis of transitional upland rice swidden cropping systems in northern Thailand. Louvain: KUL. p. 220.

PROGRAM 2.DEVELOPMENT OF INTEGRATED WATERSHED MANAGEMENT AT SCWMC IN BANDARBAN

Abstract

In hilly region of Bangladesh receive huge amount of rainfall in the country. But this rainfall was not well distributed. This huge amount of excess rainfall drains out of CHT along with eroded soil material through numerous channels (streams), canals and rivers of the hills without significant natural and artificial obstacle. Thus, following heavy downpour of the rainy season, the area suffers from severe drought and water scarcity in the dry season. Hence, it is imperative to preserve excess water of the rainy season by creating suitable dams in the hilly areas to boost up local water need round the year and secure sufficient quality water in the drier months for irrigation, fishery and other multipurpose uses. The experiment has been conducted near by the multipurpose dam at the research area of Soil Conservation and Water Management Centre, SRDI- under Bandarban Hill District. The main objectives of the research program was to manage and utilize the rain & runoff water for irrigation, fishery, poultry & household purpose, to protect, conserve and improve the slopping land of watershed, minimizing soil erosion hazard, sustained production and settled agriculture & to improve and increase the production of food, fruits, fuel and fodder and to bring scattered inhabitants in clusters to ensure healthy & safe environment. Several activities have been carried out simultaneously in suitable locations. In each activities soil conservation Technique has been introduced. Soil conservation measures of both the mechanical and bio-engineering were adopted. Mechanical measure has been accomplished with bench terracing, grassed waterways, contour trench, half moon terracing etc. Most degraded sites have already been reclaimed by using jute geo-textile. Biological method has been adopted by introducing different hedge species and which will be planted in contour lines. Fishes were introduced in the reservoirs. Irrigation for plants and seasonal crops would be done. This Research will help the hill dwellers to use the rain & runoff water for Agriculture & family life. Poultry and fish cultivation will fill-full the demand of food i.e. protein. Various food species will mitigate the demand of vitamins and minerals. It will also support them by providing cash value. Mechanical measures of soil conservation practices would be helpful for minimizing soil erosion hazard and rehabilitation of degraded lands.

INTRODUCTION

Hills of Bangladesh receive highest rainfall in the country. The average annual rainfall in CHT is 2682 mm (1941–2000; c.v. 18%) with distinct wet and dry seasons. The rainy season starts in May and continues up to October, with a sharp increase in the rainfall from May to June and the July rainfall accounting for 24% of the annual rainfall. The annual reference evapotranspiration is approx. 1350 mm leaving a potential rainfall excess of approx. 1350 mm y^{-1} . This huge amount of excess rainfall drains out of CHT along with eroded soil material through numerous channels (streams), canals and rivers of the hills without significant natural and artificial obstacle. Thus, following heavy downpour of the rainy season, the area suffers from severe drought and water scarcity in the dry season. Hence, it is imperative to preserve excess water of the rainy season by creating suitable dams in the hilly areas to boost up local water need round the year and secure sufficient quality water in the drier months for irrigation, fishery and other multipurpose uses.

Land degradation is one of the major ecological issues of the world. Land degradation means loss in the capacity of given land to support growth of useful plants on a sustained basis (Singh, 1994). Due to different types of land degradation, Bangladesh lost a substantial amount of production which in terms money may be hundreds of billion taka in every year (BARC, 1999). Faulty Jhum cultivation in hilly area causes gully erosion and loss of soil ranges from 10 to 120 t ha⁻¹yr⁻¹ (Farid *et al.*, 1992). Soil degradation is difficult to quantify and the impact of soil loss and destruction is not evident immediately. Recently, however, the magnitude of the cumulative effects has been described in some studies. Since 1970, the farmers world over have lost an estimated 480 billion tons of top soil, roughly equivalent to all of India's crop lands (Brown, 1991). In Canada, soil degradation has been costing farmers US \$ one billion a year. In India, about 173 m ha or 53% of the total geographical area are subjected to varying degrees and forms of soil erosion (Bali, 1990). Analyses of annual soil erosion rates in India have indicated that 5334 million tones (i.e. 16.33 tone/ha/year) of soil is detached annually and of this about 29% is carried away by the rivers into sea. Nearly 10% of it is being deposited in reservoirs losing 1-2% of its capacity (Narayana and Ram Babu, 1983). Scientific management of land and water resources is the key to increase productivity.

Vegetation and land use play an important role controlling the intensity and frequency of overland flow and surface erosion (Mitchell, 1990; Gafur *et al.*, 2001b). Naturally, woody perennials and tree species produce large amounts of aboveground biomass. Because of their perennial nature, there is a continuous addition of organic matter and biomass to the soil. Tree crops influence the microclimatic factors such as soil and air temperature, net radiation reaching the ground surface evaporative demand, etc. Expectedly, soil and air temperature is lower during the day in the vicinity of perennial hedges than farther away from them. Under this condition, soil organic matter content is being continuously increased, activity of soil fauna increased and soil structure improved (Lal, 1991).

So far very little scientific effort have been taken to grow different crops/annual fruits in different hill slopes of CHT, which could reduce pressure on already squeezed suitable Jhum land. Considering all these circumstances, a land use system that will ensure or sustain the production as well as conserve soil fertility and reduce soil loss should be developed particularly for the hilly regions of Bangladesh.

This proposed research program was, therefore, designed to develop technique for minimizing soil loss and fertility development to achieve better economic return along with reducing the water scarcity and ensure food security.

OBJECTIVES

- a. To manage and utilize the rain & runoff water for irrigation, fishery, poultry & household purpose.
- b. To protect, conserve and improve the sloping land of watershed, minimizing soil erosion hazard, sustained production and settled agriculture.
- c. To improve and increase the production of food, fruits, fuel and fodder and to bring scattered inhabitants in clusters to ensure healthy & safe environment.

MATERIALS AND METHODS

The experiment was conducted near by the multipurpose dam at the research area of Soil Conservation and Water Management Centre, SRDI- under Bandarban Hill District. Several activities were carried out simultaneously in suitable locations. In each activities soil conservation Technique will be introduced. Different horticultural fruit species has been planted along with hedge plant for minimizing erosion hazard. The experimental plots has been selected in such way that the area individually can be treated as a micro watershed and each plot has been considered as a treatment. The area has been divided into smaller parts depending on slope gradient and soil characteristics. Prior to plantation composite topsoil samples will collect from each site and physical, chemical and mineralogical analysis will done to compare the soil characteristics before and after cultivation. The seedlings/saplings has been planted in contour line (across the slope) with following plant spacing and row spacing on the basis of species. Preparation of land has been started at suitable time. Intercultural operations have been done as and when required. Data on yield and yield contributing characters will take at harvest time and economic analysis will made following standard statistical procedure. Soil conservation measures of both the mechanical and bio-engineering has been adopted. Mechanical measures have been accomplished with bench terracing, grassed waterways, contour trench, half moon terracing etc. Most degraded sites has already been reclaimed by using jute geo-textile. Biological method would be adopted by introducing different hedge species and which has been planted in contour lines.

Different horticultural crops like mango, Jujube (Kul) Orchard (Apple Kul & BAU Kul), Multi-fruit Garden and citrus species has been planted on adjacent to the watershed both on the flat & sloping lands following contour lines. Pits for seedlings would be dug in the month of April to May. Manures and fertilizers like cow dung, N, P & K will be applied as per recommendation. Planting of the seedlings has been done in the month of June to July depending on rain. Intercultural operation will be done manually as when and where necessary. Half moon terrace has been made to conserve moisture before the end of the monsoon. Mulch has been done in the half-moon terraced to keep the soil moisture, adding biomass & maintain the good health of the soil. Year round different seasonal vegetative crops will be grown in bench terraced area and necessary irrigation would be done from nearby multi-purposed dam water. Intercultural operations like weeding and earthling up of the soil will do equally in all treatments to get better results. Before sowing of seeds, a mixture of cowdung, N, P and K as urea, triple super phosphate and murate of potash will apply at different doses. Other management practices and preventive measures will take as and when need. Data on different attributes of different crops will record timely following standard procedures. Meteorological information's like rainfall, humidity percentage, daily maximum and minimum temperature, soil temperature in different depth will be collected.

Fishes has been introduced in the reservoirs. Experimental site also be selected near by the reservoirs. Irrigation for plants and seasonal crops would be done. Training & Demonstrations for various levels would be held on.

RESULT & DISCUSSION

Table 13. Approximate cost and return (BDT) from experimental plots

Sl. No	Particulars	Approx. Cost	Return in BDT		
			FY 2015-16	FY 2016-17	FY 2017-18
1.	Fish cultivation	93,000.00	-	20,000.00	30000.00
2.	Multi-fruit Garden (Established garden) Inter-culture only.	9,000.00	12,000.00	15,000.00	18,000.00
3.	Newly planted Mango orchard	32,000.00	Nil	500.00	1,700.00
4.	Newly planted Citrus orchard	5,000.00	1,200.00	1,500.00	1,800.00
5.	Vegetables on Bench Terrace	10,000.00	15,300.00	12,800.00	14,200.00
6.	Lichi garden (Established garden) Inter-culture only	15,000.00	60,000.00	Nil (Alternate bearing)	40,000.00
7.	Barikul Orchard (Apple Kul & BAU Kul)	1,000.00	500.00	700.00	1,400.00
8.	Coconut & battlenut (Inter-culture only)	3,000.00	3,800.00	4,200.00	4,500.00

Soil loss from degraded land under different treatment for 2016 is presented in Table 14. It was recorded that highest soil loss was recorded in control plot (15.51 t/ha) where no geo-jute or jute mat was used, followed by plot managed by 500 GSM (8.67 t/ha) and 700 GSM (5.28 t/ha). As 700 GSM jute mat is denser than 500 GSM Jute mat, so it provides better soil cover than that of 500 GSM jute mat. On the other hand 500 GSM jute mat is better than 700 GSM jute mat for growing vegetation at primary stage as it is lighter than 700 GSM Jute mat.

Table 14. Soil loss from degraded land under different treatment for 2016

Month	Soil loss (t/ha)		
	700 GSM	500 GSM	Controlled
June/2015	2.37	3.74	7.26
July/2015	1.40	2.05	3.97
August/2015	0.66	1.18	1.65
September/2015	0.43	1.05	1.36
October/2015	0.42	0.65	1.27
Total	5.28	8.67	15.51

Run-off from degraded land under different treatment for the year-2016 throughout the rainy season was calculated and shown in Table 13. It was recorded that the highest runoff was recorded in controlled plot where there was no geo-jute or jute mat, followed by plot managed with 700 GSM & 500 GSM.

Table 15. Run-off from degraded land under different treatment for the year-2016

Month	Run-off (%)			Monthly Rainfall (mm)
	700 GSM	500 GSM	Controlled	
June/2015	27.89	29.52	36.63	684
July/2015	34.61	39.41	52.85	532
August/2015	45.41	48.62	64.68	468
September/2015	29.71	32.35	35.87	359
October/2015	34.26	36.72	41.66	144

Table 16. Chemical properties of Soil under different jute-mat (Before setting the experiment)

Particulars	pH	OM%	K	Ca	Mg	TN (%)	P	S	B	Cu	Fe	Mn	Zn
		meq/100g soil					µg/g soil						
700-GSM Plot	4.5	3.36	0.36	3.43	2.08	0.40	6.70	11.3	0.20	0.76	68.8	25.2	1.50
500-GSM Plot	4.3	3.09	0.30	2.28	1.54	0.43	3.67	1.7	0.25	0.70	50.2	14.0	0.94
Control Plot	4.6	2.82	0.49	4.73	2.88	0.22	6.88	12.4	0.28	0.78	72.0	24.2	2.24

Table 17. Chemical properties of Soil under different jute-mat (After Two years)

Particulars	pH	OM%	K	Ca	Mg	TN (%)	P	S	B	Cu	Fe	Mn	Zn
		meq/100g soil					µg/g soil						
700-GSM Plot	5.9	3.77	0.39	9.08	2.71	0.19	3.28	1.11	0.03	0.69	77.0	7.23	1.48
500-GSM Plot	5.9	3.36	0.35	11.56	3.37	0.18	2.83	0.27	0.23	0.85	80.2	6.81	1.5
Control Plot	5.2	2.98	0.30	8.75	2.47	0.17	2.96	3.67	0.18	0.63	51.5	5.3	1.0

Soil nutrient are increased after setting the jute geo-textile.

CONCLUSION

This activity has been proposed aiming to mitigate the basic need of a small family consisting of five members. This Research will help the hill dwellers to use the rain & runoff water for Agriculture & family life. Poultry and fish cultivation will fill-full the demand of food i.e. protein. Various food species will mitigate the demand of vitamins and minerals. It will also support them by providing cash value. Other crops like coconut and battle-nut will enrich the economy in various ways. Mechanical measures of soil conservation practices would be helpful for minimizing soil erosion hazard and rehabilitation of degraded lands. Seasonal vegetables grown in Bench Terrace will supply them food as it is easy to irrigate on bench terraced area for suitable crop's yield. Other mechanical measures i.e. half moon terracing would be helpful to conserve moisture for the drought time and added mulching would enrich the soil fertility by increasing organic matter in soil. Barikul (BARI kul and Apple kul) is a naturally drought tolerant plant and suitable for drought prone areas. It has also great demand in local market. Irrigation for different orchard (Mangoes, litchi, malta, lemon etc) can be done from the nearby multi-purpose reservoir. On the other hand, the reservoir can serve the purpose water used for house hold activities of the hill dwellers. The economical return from the experiment will be obtained & calculated for the next time. Further the whole activities would be used for training and demonstration purposes.

REFERENCES

- Bali, J.S. (1990). Integrated Watershed Management- A National Perspective- Proc. International Symp. on Water erosion, Sedimentation and Resource Conservation (Oct. 9-13, 1990), 279-292.
- BARC. 1999. Land degradation situation of Bangladesh. Soil Science Division. Bangladesh Agricultural Research Council, Farmgate, Dhaka.

- Brammer, H. 1986. Reconnaissance soil and land use survey: Chittagong hill tracts (1964-65), Soil Resource Development Institute, Dhaka. pp. 98.
- Brown, A.G. 1988. Soil development and geomorphic processes in a Chaparral Watershed: Rattlesnake Canyon, S. California, USA. *Catena Suppl.* 12: 45-58.
- Farid, A. T. M., A. Iqbal, and Z. Karim. 1992. Soil erosion in the Chittagong hill tract and its impact on nutrient status of soil. *Bangladesh J. Soil Sc.* 23 (122): 92-101.
- Gafur, A., J.R. Jensen, O.K. Borggaard and L. Petersen. 2001b. Runoff and losses of soil and nutrients from small watersheds under shifting cultivation (Jhum) in the Chittagong Hill Tracts of Bangladesh. *J. Hydrol.*
- Gafur, A., O.K. Borggaard and L. Petersen. 2001c. Economic appraisal of shifting cultivation in the Chittagong Hill Tracts of Bangladesh. *Ambio* submitted.
- Khybri, M.L. 1983. Soil erosion problems and soil conservation in India. *Watershed management*. Bhatt. P.N. (ed.). pp. 6-13.
- Singh, H.B. 1994. Operational Research Projects of Watershed Management in Gujrat. (Personal communication).
- Shoaib. 2000. Development of sustainable cultivation practices for minimizing soil erosion on hill slope. Soil Resource Development Institute (SRDI).
- Khisha, S.K., J.U.M. Shoaib and N.A. Khan. 2002. The experience of promoting slope agricultural land technology for hillside farms in the Chittagong Hill Tracts. In: "Farming Practices and Sustainable Development in the Chittagong Hill Tract" (N.A. Khan *et al.*, Eds.). CHTDB and VFFP, Intercooperation, pp. 114-115.
- Khisa, S. Kanti, J.U.M. Shoaib, and K.N. Ahmed. 2006. Selected natural resources conservation approaches and technologies in Chittagong Hill Tracts Bangladesh. *Bangladesh Conservation Approaches and Technology (BANCAT)*.
- Khybri, M.L. 1983. Soil erosion problems and soil conservation in India. *Watershed management*. Bhatt. P.N. (ed.). pp. 6-13.
- Mitchell, D.J. 1990. The use of vegetation and land use parameters in modelling catchment sediment yields. In : J.B. Thornes (Ed.). *Vegetation and Erosion, Processes and Environments*. Wiley, Chichester, pp. 289-314.
- Singh, G.S., G. Shastri and S.P. Bhardwaj. 1990. Watershed response to conservation measures under different agro-climatic regions of India. *Indian J. Soil Conservation*. 18 (3): 16-22.

PROGRAM 3. STUDY ON MANAGEMENT AND ECONOMIC VALUE OF *SCHUMANNIANTHUS DICHOTOMA* (MURTA/ PATIBET) IN HILLY JHIRI LAND AT BANDARBAN

Abstract

Schumannianthus dichotoma (Murta/Patibet) is widely grown in wetland areas of Bangladesh, providing the raw materials for prayer and bed mats and also minimizing soil erosion. A suitable field situated in a Jhiri locating of the South-south-east side of the SCWMC administrative Building has been selected for cultivation of *Patibet*. The main objectives of the research program was to study the suitability and yield or productivity of Murta in Hilly Region of Bangladesh, to ensure the fallow lands of hilly Jhiri in to productive and minimize soil erosion hazard & to strengthen the economical efforts of the hill dwellers by increasing off farm activities & to supplement the traditional Jhum Practices. Propagation was from rhizomes and branch cutting, and little intensive management was required. Harvesting was usually done annually, from mid September to the end of March. Soil has been made up for proper propagation. Necessary intercultural operation is going on accordingly. Weeding is needed only for vines and climbing weeds, generally before the rainy season. Weeding, especially of the main weed, Asam lata (*Eupatorium odoratum*), should usually be carried on along with harvesting, or occasionally. There were no significant pest and disease attacks in *Patibet* plantations.

Patibet can play a vital role in the economy and environment CHT of Bangladesh. It can easily be cultivated in hilly Channel/Jhiri that remain fallow and remain wet even in the dry season. These lands are not suitable for cultivation of other cash crops. The cultivation of *Patibet* is inexpensive and does not conflict with the production of agricultural crops. This program will minimize soil erosion hazard in Chittagong Hill Tracts. This study will ensure income generating crops instead of harmful jhum cultivation and safe hill environment.

PROGRAM 4. EFFECT OF DIFFERENT HEDGE SPECIES ON CONTROLLING SOIL EROSION, RUNOFF, AND NUTRIENT MINING OF SNACK GOURD AT GENTLE SLOPE IN BANDARBAN

Abstract

The study involving Snack Gourd conducted at the Soil Conservation and Watershed Management Centre (SCWMC), Soil Resource Development Institute (SRDI), Bandarban to investigate the effect of Effect of different hedge species and slope gradient on controlling soil erosion, runoff and nutrient mining at gentle slope in CHT. The treatments were: T₁: *Indigofera* hedge, T₂: Pineapple hedge and T₃: Control (without hedge). Slope percent of the experimental plot was 12%. Hedge species were planted in following contour lines maintaining 5 m alley distance. Measurement of soil loss and run-off was carried out by established and locally fabricated multi-slot divisors. Natural condition of the slopes was not disturbed or no any soil work was done to make artificial slope. Species of hedge plants have a great effect on plant growth and crop yield. Hedge plant of low height (*pineapple*) provides a better performance than that of higher height hedge plant because it provides intensive light & better root binding. But higher height hedge plant provides more bio-mass than lower height hedge plants. Performance of pineapple hedge species on soil loss minimizing capacity was recorded the best on all slope

gradients. Hedge always plays a vital role on plant growth, crops productivity, fruit length & weight as well as minimizing of soil erosion.

INTRODUCTION

The conservation of soil and water is essential for sustainable production, environment preservation and balanced ecosystem (Sarma *et al.* 2000). Loss of soil by water erosion on sloping lands adversely affects the physical, chemical and biological properties of soils, leading to low crop productivity (Larson *et al.* 1985 and Sur *et al.* 1994). Land use change associates erosion is mostly responsible for land degradation and desertification in different part of Asia and Africa, bringing about large reduction in vegetation growth, siltation of water courses, filling of valleys and reservoirs and the formation of deltas along the coastal areas. Erosion is accompanied by deposition of alluvial materials by flooding and filling of valleys, waterways or extending coastal plains and deltas towards the sea. Contour hedgerows are also effective in controlling run off and soil erosion and improve soil physical properties. Control plots have higher run off and soil loss than those plots with hedgerow (Khisa *et al.* 1999). Uddin and Firoz (2001) recommended hedgerow for cultivation on sloppy land. They described that hill slope may be divided into a series of alley separated by hedgerow on contour lines, because hedgerow plants are effective in controlling soil erosion and reducing run off. A number of research works have been conducted in the tropics regarding soil fertility improvement under agroforestry practices. Agboola *et al.* (1982) as cited by Attah-Krah and Sumberg (1988) reported that the soil chemical properties like pH, OM, N, available P and CEC improved with the use of *Gliricidia sepium* as hedgerow. Attah-Krah *et al.*, (1986) as cited by Nair (1993) reported that organic matter content and nutrient levels of soil were higher under alley cropping as compared to soil without trees.

Trees and shrubs have several functions to control erosion like (i) increase soil cover, by litter and pruning (ii) provide partly permeable hedgerow barriers (iii) lead to the progressive development of terraces, through soil accumulation upslope of hedgerows (iii) increase soil resistance to erosion, by maintenance of organic matter (iv) stabilize earth structures by root systems and (v) make productive use of the land occupied by the conservation works (Young 1989 a). Alley cropping or hedgerow cultivation is very helpful in controlling of soil erosion in the hilly area. Hill Tract Development Board of Bangladesh identified five nitrogen fixing tree species like *Leucaena leucocephala*, *Gliricidia sepium*, *Indigofera tysimani*, *Flemingia spp*, and *Desmodium rensonii* etc. and two grass species *Vetiviera zizanoides* and *Thysanolaena maxima* for controlling runoff and erosion in the hilly region of Bangladesh (Khisa *et al.*, 2002). Singh *et al.* (1990) found that runoff and soil loss were substantially reduced when small watersheds with agriculture were replaced either by trees and grasses (silvipasture) or with mechanical measures. In a study, Wiersum (1984) found that different agroforestry systems cause lowest soil erosion. Facing the location specific, environmental friendly agricultural development challenges in CHT, Soil Conservation and Watershed Management Center (SCWMC), Bandarban has stepped up its efforts to generate scientific information on the major land use practices in the region, This study is the first attempt to investigate and quantify the effect of different hedge species on soil loss and run-off and its economic performance on crop cultivation. Therefore, the present study was undertaken to find out the feasible hedge species for controlling soil erosion and for crop cultivation in the CHTs.

OBJECTIVES

- a. To compare soil loss, runoff and nutrient mining in cultivating Snack Gourd under different hedge species.
- b. To evaluate the economical aspects of cultivation under different hedge species.

MATERIALS AND METHODS

The experiment was carried out under non replicated condition. Three experimental plots of 100 sqm (5 m x 20 m) on gently (12%) were selected in the SCWMC, Bandarban. Two different species were used as hedge species like; *Indigofera* and *Pineapple*. Hedge species were planted in 2013 following contour lines maintaining 5 m alley distance but the data of the experimentation were recorded during 2016. Measurement of soil loss and run-off was carried out by established and locally fabricated multi-slot divisors. Natural condition of the slopes was not disturbed or no any soil work was done to make artificial slope. Pits were dug 1.80 m c/c in rows maintaining contour. Total number of pits in each plot was 24. Fertilizers were applied as per recommendation of soil test value. 3 to 4 seeds were sown in each pit.

After germination, two seedlings were allowed for yield. Cultural operations were done as usual in all the plots. Soil loss and run-off from each 100sqm (5m x 20m) experimental plots were measured after each shower throughout the rainy season. Daily and eventually monthly soil loss and run-off were estimated from each treatment by processing aliquot of sample every day. Every morning (if rains previous day) amount of run-off water is measured in the multi-slot and aliquot of about 2 liter is sampled from each tank. Suspended sediment in the sampled aliquot is measured by simple filtering and oven drying. Corresponding rainfall is recorded from the automatic and ordinary rain gauge of SCWMC. Climatic data like rainfall, temperature, humidity, evaporation etc. were recorded daily. Snack Gourd was selected as a test crop. Different agronomic practices were done when it was necessary.

RESULTS AND DISCUSSION

Soil loss under different hedge species in gentle slope at Snack Gourd growing plot during 2016-18 sessions is presented in table 23. Soil loss under Snack Gourd at the alley of different hedge species throughout the rainy season was calculated. It was recorded that highest soil loss was recorded in control plot (15.35 &16.04 t/ha) where no hedge species were used, followed by *indigofera* hedge species used plot (8.57 &6.06 t/ha) and pineapple hedge species used plot (5.05 &5.46t/ha).

Table 18. Soil loss under the cultivation of different hedge species in 2016-17 (t/ha/year)

Particulars	Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Total
Indigofera	2016-17	-	-	-	-	-	2.45	3.09	1.34	0.62	1.07	-	-	8.57
	2017-18	-	-	-	-	0.85	1.94	2.57	1.45	1.63	0.62	-	-	9.06
Pineapple	2016-17	-	-	-	-	-	1.19	1.80	0.69	0.41	0.96	-	-	5.05
	2017-18	-	-	-	-	0.61	0.85	1.78	0.84	1.01	0.37	-	-	5.46
Control	2016-17	-	-	-	-	-	3.85	6.04	2.66	1.09	1.71	-	-	15.35

	2017-18	-	-	-	-	1.83	3.36	3.81	1.73	3.68	1.63	-	-	16.04
--	---------	---	---	---	---	------	------	------	------	------	------	---	---	-------




Figure 43. Soil loss (t/ha/year) under the cultivation of different hedge species in 2016-18

Runoff under Snack Gourd at the alley of different hedge species throughout the rainy season were calculated (Table 24). It was recorded that highest runoff was recorded in control plot where no hedge species were used, followed by *indigofera* hedge species used plot and pineapple hedge species used plot.

Table 19. Run off under the cultivation of different hedge species in 2016-2018

Particulars		Month wise runoff (%) 2016-18											
		Jan	Feb	Mar	Apr	May	June	July	Aug	Sep	Oct	Nov	Dec
Indigofera	2016-17	-	-	-	-	-	43.14	52.79	49.45	35.98	28.23	-	-
	2017-18	-	-	-	-	7.07	41.14	28.41	19.21	25.30	22.41	-	-
Pineapple	2016-17	-	-	-	-	-	36.57	49.80	47.25	32.46	25.76	-	-
	2017-18	-	-	-	-	6.39	36.69	26.32	18.17	17.19	16.85	-	-
Control	2016-17	-	-	-	-	-	46.11	56.22	51.66	39.49	33.17	-	-
	2017-18	-	-	-	-	9.77	44.79	33.64	22.34	31.29	29.80	-	-

CONCLUSION

1. Use of different hedge has created positive effect on the morphological and reproductive characteristics as well as at the yield of Snack Gourd.
2. Between two hedges pineapple hedge reduced relatively higher soil loss due to its soil binding capacity of roots.
3. Between two hedges pineapple hedge reduced relatively higher runoff.

REFERENCES

Annual Report, 2010. Soil Conservation and Watershed Management Centre, SRDI, Bandarban. pp.

- Khisa, S.K. 2001. Contour hedgerow inter-cropping agroforestry technology for degraded hill side farms in Chittagong hill tracts. Proc. National Workshop on agroforestry research, Gazipur, September 16-17. pp. 179-184.
- Khisa, S.K.; M.K. Alam and N.A. Siddiqi. 1999. Broom grass (*Thyanolaena maxima*) Hedgerow : A Bioengineering Device for Erosion and Slope Stabilization. Proc. First Asia-Pacific Conference on Ground and Water Bioengineering for Erosion and Slope stabilization, April 19-21, 199, Manila, The Philippines International Erosion Control Association. pp. 172-177.
- Khisha, S.K., J.U.M. Shoaib and N.A. Khan. 2002. The experience of promoting slope agricultural land technology for hillside farms in the Chittagong Hill Tracts. In : "Farming Practices and Sustainable Development in the Chittagong Hill Tract" (N.A. Khan *et al.*, Eds.). CHTDB and VFFP, Intercooperation , pp. 114-115.
- Khisa, S. Kanti, J.U.M. Shoaib, and K.N. Ahmed. 2006. Selected natural resources conservation approaches and technologies in Chittagong Hill Tracts Bangladesh. Bangladesh Conservation Approaches and Technology (BANCAT).
- Khybri, M.L. 1983. Soil erosion problems and soil conservation in India. Watershed management. Bhatt. P.N. (ed.). pp. 6-13.
- Lal,R. 1993. No-tillage farming. Soil and water conservation and management in the humid and sub-humid tropics. iiTA Tech. Monograph2:64.
- Singh, G.S., G. Shastri and S.P. Bhardwaj. 1990. Watershed response to conservation measures under different agro-climatic regions of India. Indian J. Soil Conservation. 18 (3): 16-22.
- Shoaib. 2000. Development of sustainable cultivation practices for minimizing soil erosion on hill slope. Soil Resource Development Institute (SRDI).
- Uddin, M.S. and Z.A. Firoz. 2001. Yield of lemon and guava as influenced by hill slope and hedgerow plants. Proc. National workshop on agroforestry research, Gazipur, September 16-17. pp. 196-198.

PROGRAM 5. STUDYING SOIL LOSS AND YIELD PERFORMANCE OF PINEAPPLE BASED JACKFRUIT ORCHARD ON HILL SLOPE FOLLOWING CONTOUR LINE

Abstract

The study was conducted at the Soil Conservation and Watershed Management Centre (SCWMC), Soil Resource Development Institute (SRDI), Bandarban .Area of the selected site is 61.0 m x 21.0 m (200'-0" x 70'-0") likely about 0.32 acre. Average slope of the site was 35%. The main objectives of the research program were to introduce technique for effective land use in achieving food security as an alternative farming system, to study the yield of pineapple as an intermediate crop with permanent horticulture & to determine soil loss. Contour lines were selected at 0.50 meter vertical interval. Pineapple suckers were planted in the predefined contour with application of necessary manures and fertilizers. Pits for planting Jackfruit were dug before

plantation of pineapples' sucker. Soil loss was recorded in peg methods. 21 number Color pegs were inserted in to a certain depth at a distance of 10.0 m. peg to peg and row to row before the rain. Intercultural operation is going on as necessary. Jackfruit was main horticultural crop and pineapple was intermediate crops. Main crops will need a certain period for its production. Pineapple was introduced as an intermediate crop for introducing technique of effective land use for achieving food security and also an alternative farming system. The intermediate crop has been planted across the slope following contour. It will act as hedge rows which will be very helpful for minimizing soil erosion hazard as well as providing for the intermediate period.

INTRODUCTION

Land, water and vegetation are the most important natural resources for providing environmental and livelihood security to the mankind. They provide food, fodder, firewood, fibre and other material needs of the people. History bears testimony to the high regard that man holds for these resources. However, with the advent of civilization leading to cultivation of land and subsequently ever increasing pressure of man and animal population, the natural balance between these resources has been distorted and as a consequence, serious problems of soil and water conservation have arisen. The movement of water on land if not properly managed may cause soil erosion and render land incapable of sustained production. Land degradation is one of the major ecological issues of the world. Land degradation means loss in the capacity of given land to support growth of useful plants on a sustained basis (Singh, 1994). Due to different types of land degradation, Bangladesh lost a substantial amount of production which in terms money may be hundreds of billion taka in every year (BARC, 1999). Faulty Jhum cultivation in hilly area causes gully erosion and loss of soil ranges from 10 to 120 t ha⁻¹yr⁻¹ (Farid *et al.*, 1992). Soil degradation is difficult to quantify and the impact of soil loss and destruction is not evident immediately. Recently, however, the magnitude of the cumulative effects has been described in some studies. Since 1970, the farmers world over have lost an estimated 480 billion tons of top soil, roughly equivalent to all of India's crop lands (Brown, 1991). In Canada, soil degradation has been costing farmers US \$ one billion a year. In India, about 173 m ha or 53% of the total geographical area are subjected to varying degrees and forms of soil erosion (Bali, 1990). Analyses of annual soil erosion rates in India have indicated that 5334 million tones (i.e. 16.33 tone/ha/year) of soil is detached annually and of this about 29% is carried away by the rivers into sea. Nearly 10% of it is being deposited in reservoirs losing 1-2% of its capacity (Narayana and Ram Babu, 1983). Scientific management of land and water resources is the key to increase productivity.

Soil erosion in agricultural systems is a very important to manage. If the productive layer or topsoil is eroded away, then the ground is very unproductive in producing crops. Soil erosion generally occurs only on slopes, and its severity increases with the severity of slope. The Chittagong Hill Tracts represents a very fragile hill ecosystem and is characterized by steep to extremely steep slopes with ninety percent of its landscape belonging to upland category that limits its land capabilities. Most of the slopping areas are closely dissected and sharp ridged, slopes are mostly steep to very steep dominated by shallow to moderately deep, nutrient poor, loamy to clayey and slightly to very strongly acid soils. These factors combined cause soil erosion, siltation of lakes and rivers and soil fertility decline thereby creating a food insecurity situation in the region. Major agricultural activity in this area is traditional rainfed farming which

is locally known as 'Jhum' and commonly known as 'Shifting cultivation' or 'Slash and burn' farming system. About 1.0 million peoples in CHT of which 13 different ethnic groups are directly or indirectly depend on jhum (Shoaib, 2000). It is estimated that it takes 300-1000 years to form an inch of soil. In areas of seriously erosion, this one inch of soil may be lost in a couple of years (Khybri, 1983). Jhum, the dominant form of land use in CHT, widely practiced by tribal communities and remains as a major source of livelihood for most of the hill people. It is estimated that about 26,000 families practices Jhum every year and about 1,43,000 people depend on Jhum for subsistence. This cultivation system has become unsustainable because diminished suitable land availability has forced the tribal communities into shorter and shorter jhum cycles (now down to 3-5 years per cycle) there by reducing soil fertility and increasing soil erosion hazards. Agroforestry has been considered to be financially more attractive than jhum. Environmentally it also appeared to be more suitable for CHT as the rate of soil erosion under this system was found considerably lower than under other land use systems such as jhum and root crops (Gafur, 2001).

Vegetation and land use play an important role controlling the intensity and frequency of overland flow and surface erosion (Mitchell, 1990; Gafur *et al.*, 2001b). Naturally, woody perennials and tree species produce large amounts of aboveground biomass. Because of their perennial nature, there is a continuous addition of organic matter and biomass to the soil. Tree crops influence the microclimatic factors such as soil and air temperature, net radiation reaching the ground surface evaporative demand, etc. Expectedly, soil and air temperature is lower during the day in the vicinity of perennial hedges than farther away from them. Under this condition, soil organic matter content is being continuously increased, activity of soil fauna increased and soil structure improved (Lal, 1991).

So far very little scientific effort has been taken to study the yield of short duration crop (pineapple) as an intermediate crop with permanent horticulture which could reduce pressure on already squeezed suitable Jhum land. Considering all these circumstances, a land use system that will ensure or sustain the production as well as conserve soil fertility and reduce soil loss should be developed particularly for the hilly regions of Bangladesh. This proposed research program was, therefore, designed to develop technique for minimizing soil loss and fertility development to achieve better economic return and ensure food security.

Objectives

1. To introduce technique for effective land use in achieving food security as an alternative farming system.
2. To study the yield of pineapple as an intermediate crop with permanent horticulture.
3. To determine soil loss in peg method.

MATERIALS AND METHODS

The research has been introduced in the financial year 2016-2017. Site was selected to the east side of the farm shed at SCWMC Research station, SRDI, Bandarban. Area of the selected site is 61.0 m x 21.0 m (200'-0" x 70'-0") likely about 0.32 acre. Average slope of the site is 35%. After selection of the plot, the jungle was cleaned by manual labour. Contour lines were selected at 0.50 meter vertical interval. Pineapple suckers were planted in the predefined contour with application of necessary manures and fertilizers. Pits for planting Jackfruit were dug before

plantation of pineapples' sucker. Recommended dose of fertilizers and manure were applied in to the pits. Pineapple suckers were collected and planted in May 2016. There are 14 rows of pine apple plantation. Saplings of Jackfruit were collected from horticulture centre. Jackfruit's saplings were planted in the pits maintaining plant to plant and row to row distance 25'-0" in July 2016. There are 18 nos. of jackfruit sapling were planted in this plot. 21 number Color pegs were inserted in to a certain depth at a distance of 10.0 m. peg to peg and row to row before the rain. The pegs were 1'-6" long and were colored by red & white every 6". One third length of the peg were driven in the soil and two third were above the soil. Grounds RL near different pegs were recorded before the rain and after rain in each year. The Cumulative difference of the two depths is considered as the depth of transported soil. Intercultural operations were done manually when necessary. Yield performance of pineapple is being studied. It was found that near about 50% of the pineapple plans are on bearing stage. The pineapples will be destroyed when the jack fruits are being harvested.

RESULTS AND DISCUSSION

Soil loss by peg method was studied for 2016-2017 and 2017-2018. There were three rows of peg in the plot in combination of seven pegs in each row. At first soil loss for seven pegs of upper, middle and bottom rows were determined by using leveling instrument. Then the average depth of transported for each row were calculated accordingly. It was found that the average depth of transported soil were 0.655 mm, 0.710 mm and 0.780 mm for the pegs of upper, middle and bottom lines respectively for 2016-2018. On the other hand, the average depth of transported soil for upper, middle and lower parts were 0.60 mm, 0.62 mm and 0.67 mm respectively during the period of 2017-2018. Then the average depths of soil loss for three rows were calculated and it was found 0.715 mm and 0.630 mm for the year 2016-2017 and 2017-2018 respectively. Finally the total soil loss soil loss was calculated assuming 1mm depth soil loss is equal to 13.70 t/ha and it was 9.796 t/ha and 8.631 t/ha per year. It was also found that both the main crop and the intermediate crops were growing satisfactory. A data for determination of soil loss is given below:

Table 20. Soil Loss under pineapple based Jackfruit orchard on steep slope

Location of peg	Average soil loss in mm		Average of all rows (mm)		Total soil loss (t/ha)		Mean total soil loss (t/ha)
	2016-2017	2017-2018	2016-2017	2017-2018	2016-2017	2017-2018	
Upper row	0.655	0.60	0.715	0.63	9.796	8.631	9.212
Middle row	0.710	0.62					
Bottom row	0.780	0.67					

Table 21. Yield and return from pineapple based Jackfruit orchard on steep slope

Financial year	Yield (main and associated crop)		Economical return		Remarks
	Pineapple	Jack fruit	Pineapple	Jack fruit	
2016-2017	Initial stage	Initial stage	Initial stage	Initial stage	
2017-2018	Flowering stage	Growing stage	1000.00 (Expected)	Growing stage	

CONCLUSION AND RECOMMENDATION

The conductive research is being carried on. Here, Jackfruit is main horticultural crop and pineapple is intermediate crops. Main crops need a certain period for its production. Pineapple was introduced as an intermediate crop for introducing technique of effective land use for achieving food security and also an alternative farming system. The crops are in growing stage. In the mean time, the both crops are going to be matured for its production. It is found that near about 50% plants of the intermediate crops are on bearing stage in the month of May-2018 whereas only one tree of the main crop is bearing only one jackfruit. It is also expected that Tk. 1,000/- could be earned by selling the pineapples in the local market and earning from main crop is negligible. Soil loss data for the year 2016-2017 and 2017-2018 has been recorded accordingly and it was found on average 9.796 ton $\text{ha}^{-1}/\text{year}$ and 8.631 ton $\text{ha}^{-1}/\text{year}$ respectively. The intermediate crop has been planted across the slope following contour. It will act as hedge rows which will be very helpful for minimizing soil erosion hazard as well as providing for the intermediate period. Intercultural operation is going on as necessary.

REFERENCES

- DANIDA. 2000. Interim Planning Mission, Watershed Development Project, Chittagong Hill Tracts (CHT), Bangladesh (draft).
- Gafur, A.,J.R. Jensen, O.K. Borggaard and L. Petersen.2001b.Runoff and losses of soil and nutrients from small watersheds under shifting cultivation (Jhum) in the Chittagong Hill Tracts of Bangladesh. J. Hydrol.(Not yet published).
- Khisa; S.Kanti, Shoaib, M. Jalaluddin and K.N.Ahmed.2006.Selected natural resources conservation approaches and technologies in Chittagong Hill Tracts Bangladesh. Bangladesh Conservation Approaches and Technology (BANCAT)
- Lal, R. 1983. No-tillage farming. Soil and water conservation and management in the humid and sub-humid tropics-ii TA Tech.Monograph2.64.
- Mitchell, D. J. 1990.The use of vegetation and land use parameters in modeling catchment sediment yields. In: J.B.Thornes (Ed.).Vegetation and Erosion, Processes and Environments.Wiley,Chichester,pp.289-314.
- Soil Conservation and Watershed Management Center (SCWMC), 2004. Annual Report, Soil Conservation and Watershed Management Centre, SRDI,Bandarban, Pp.7-8.
- Shoaib.2000.Development of sustainable cultivation practices for minimizing soil erosion on hill slope. Soil Resource Development Institute (SRDI).
- Soil Resources Development Institute(SRDI).1986. Reconnaissance Soil and Land Use Survey: Chittagong Hill Tracts. Forestal, Canada 1964-1965.Soil Resources Development Institute, Dhaka.206 p.
- Khybri, M. L .1983. Soil erosion problems and soil conservation in India. Watershed management. Bhatt. P. N. (ed.).pp.6-13.

PROGRAM 1. INTRODUCTION OF BENCH TERRACE FOR DEMONSTRATION AND YEAR ROUND CROP PRODUCTION

OBJECTIVES

- To reduce the quantum of overland flow/sheet flow or runoff, and their velocity.
- To minimize the soil erosion.
- To conserve soil moisture.
- To conserve soil fertility and to facilitate farming operations on sloping land.
- To promote intensive land use, permanent agriculture and checking shifting cultivation on steep lands.

MATERIALS AND METHODS

A suitable site for bench terrace was being selected at Faruk para (Chhaingya Marma para) for demonstration. It is situated just south of the Nilachal Parjaton Area and just down the Tiger Hill. Actually the site was selected to a farmer's field. The spot area of the Bench Terrace was nearly 1.0 acre & the slope gradient of the spot was nearly gentle to steep (Average- 26%). There are 11 number of Bench having variable width (2.75m to 1.0m) with a vertical interval of 0.60m. The bench terrace is made mainly inward typed and the risers are constructed maintaining 1:1 slope. The risers are protected against erosion by putting grass sod. The top portion of the spot was formed very wide and broad based terrace where tillage operation could be operated using modern tools without any difficulties. Various Robi & kharif crops like Red Amaranthus, Mustard, Bean, sweet Gourd, Pumpkin, Radish, Felon, Khaishya, kalmi, Spinash, tomato, cabbage, cauliflower etc were grown there successfully.

CONCLUSION

Terracing is one of the most accepted measures of controlling erosion on sloping and undulated lands. It is widely adopted in many countries of the world. Suitable bench terrace will facilitate intensive cultivation make the land suitable for multiple use in hilly areas. It is also helpful to increase the beauty of the land along with increasing the stability. At present most of the farmers are practicing Jhum on hill slope, which accelerate erosion. On sloping lands farmers usually can't use fertilizer or other input to produce more crops per unit of land. Bench terrace helps in proper water management, application of fertilizers or manure. It will also help to increase cropping intensity within a stable farming system. Now a day, Bench Terrace is widely being used in the hilly areas of India, Nepal, Srilanka, Tamilnadu etc. But the hill dwellers are not concerned about the construction, use and benefit of the Bench Terrace. Considering above factors study of sustainability of Bench terrace has been taken account in the farmer's field. As the measure is very cost effective, so widely subsidy is very essential to popularize the Bench Terracing mainly in the hilly areas in our country.

PROGRAM 2. REHABILITATION OF DEGRADED/ERODED SLOPING LAND BY JUTE GEO-TEXTILE ON DIFFERENT HILL SLOPES OF CHT

OBJECTIVES:

- 1) To study the effectiveness of geo-jute (untreated) in controlling soil erosion.
- 2) To rehabilitate degraded/eroded/landslide hilly areas
- 3) To stabilize/rejuvenate degraded/landslide areas of CHT

MATERIALS AND METHODS

Several degraded land at different places like Raisha Para, Satkomol para, etc. were chosen to rehabilitate by using jute geo-textile.

1. Locally available grass mulches were given on the selected degraded surface.
2. Geo-jute had been spread over the area under treatment with some exposure at grass or tree planting points.
3. The two adjoining width were generally overlapped by about 10 cm and fastened with jute threads.
4. Wooden sticks were inserted into the ground throughout geo-jute strand so that the geo-jute touches the earth surface.
5. Grasses like creeper (*Mimosa invisa*), broom grass (*Thysanolaena maxima*), khagra (*Saccharum spontaneum*) etc. were planted to check erosion by spreading soil cover and root zone in exposed locus on geo-jute strands at close spacing.
6. 50-100 pits per 100 m² are made to put seeds of different grasses & live plants.
7. Quick growing shrubs/ trees were also be planted in pits for root zone netting.

Depending upon the condition of the field, 1 to 3 nos. of drains/grass waterways were established at the upper end to check runoff water flash into the landslide area that may cause further extension of the divested area in either.

No. of Implementation Areas: 4 (Four)

CONCLUSION

1. Like any other natural fibre, jute geo-textile gets biodegraded in soil.
2. The live plants will give vegetation coverage and soil losses will be minimized at the area treated with jute geo-textile.
3. The decomposition of the fibre will take place within the ecological process that assists in the retention of moisture, improvement of soil permeability and establishment of vegetation.

PROGRAM 3. ESTABLISHMENT OF DIFFERENT HEDGE SPECIES IN FARMERS' FIELD AS TRANSFERABLE TECHNOLOGY IN CHT

OBJECTIVES

- a) To introduce modern hill cultivation and suitable technology for Soil Conservation and Watershed Management.
- b) To mitigate the need of fuel, fodder and economical purpose of the hill dwellers.
- c) To minimize soil erosion hazard.
- d) To increase bio-mass in soil properties.
- e) To accelerate the infiltration and water holding capacity of soil.

MATERIALS AND METHODS

The experiments were conducted at farmers' field in different areas like Gonespara, Meghla, Talukdar para, Parjatan Chakmapara, Raicha, Balaghata para and Nilachol etc. Different hedge species like Pine apple (fruit), Bogamedula (Shrubs), Nappier & Vetivera, (Grass), Broom, Khagra (local economical crop) etc. were selected depending on the slope, crops, need and demands of the beneficiaries. The sites may be of either newly planted or preparing for plantation. It may also be agricultural land if the field slope is gently to moderately slopping. Firstly the sites were cleaned. Then contour lines having interval of 4-6m horizontally were fixed depending on the used fruit species and the expectation of the beneficiaries by using leveling instrument or field made most temporarily use 'A-frame'. Hedge was practiced in single line. After fixation of spacing of hedge line, the hedge areas were cleaned properly. Then organic and inorganic fertilizer was mixed in the slightly dibbled pits. Seeds/sucker/ seedlings of the selected hedge was sown/planted in those pits maintaining contour lines. Gap filling was made sure if and when necessary. Intercultural operation had also been carried out. When the plants' height of the hedge exceeded 70 cm, the hedge plants were cut down at knee height level. After cutting the hedge plants, the residues part of the hedge were spread just in front of the hedge plants for filtering/arresting the eroded soil from the upper ends.

No. of Implementation Areas: 10 (ten) sites

CONCLUSION

The conservation of soil and water is essential for sustainable production, environment preservation and balanced ecosystem. Loss of soil by water erosion on slopping lands adversely affects the physical, chemical and biological properties of soils, leading to low crop productivity. Contour hedgerows are also effective in controlling run off and soil erosion and improve soil physical properties. Controlled plots have higher run off and soil loss than those plots with hedge row. Hill slope may be divided into a series of alley separated by hedgerow on contour lines, because hedgerow plants are effective in controlling soil erosion and reducing run off.

Trees and shrubs have several functions to control erosion like (i) increase soil cover, by litter and pruning (ii) provide partly permeable hedgerow barriers (iii) lead to the progressive development of terraces, through soil accumulation upslope of hedgerows (iii) increase soil resistance to erosion, by maintenance of organic matter (iv) stabilize earth structures by root systems and (v) make productive use of the land occupied by the conservation works . This study was, therefore, designed to select suitable hedge species and their alley width in respect to slope which minimized soil loss and increase crop yield.

PROPOSED RESEARCH PROGRAM (2018-2019)

Program 1. Effect of indigenous & modern cultivation methods of pineapple on soil erosion, runoff, nutrient mining at CHT

Objectives

1. To estimate & compare soil loss, runoff and nutrient mining under indigenous and modern cultivation systems of pineapple.
2. To calculate effect of soil loss on soil chemical properties.
3. To create awareness about soil conservation & watershed management among hill dwellers.

Expected output/justification:

The conservation of soil and water is essential for sustainable production, environment preservation and balanced eco system. Loss of soil by water erosion on sloping lands adversely affects the physical, chemical and biological properties of soils, leading to low crop productivity, use of indigenous methods of pineapple cultivation has created negative effect on soil erosion. This program improves soil healthy by reducing soil erosion hazard and enhances crop production. This research his encourages the hill dwellers to adopt the cultivation system to losing zero tillage across the hill slope in contour.

Program 2. Effect of plantation of Bamboo for erosion control and its economic purposes

Objective:

1. Reclamation of gullied land by minimizing erosion hazards.
2. For Landscape, aesthetic and economic purposes.
3. For environmental and ecological conservation.

Expected output:

Bamboo has evergreen leaves, dense canopy and numeral culms which can help to intercept considerable amount of rainfall. Falling raindrops change their direction and ways and reduce velocity, and therefore decrease soil erosion after multiple interceptions by tens of shoot layers and larger amount of culms. They respond to the variability in precipitation by alternating their plan form, channel geometry and sedimentation pattern which in turn influence floods and associated landscape evolutions (Rao *et al.* 2012).

Bamboo can improve poor soil conditions by building up humus content, increasing beneficial microbiological activity and improve soil structure. Bamboo leaves can filter air pollutants, recycle CO₂ and replenish the atmosphere with Oxygen. It helps to purify our streams/water supplies by taking up excess nutrients. Bamboo is also helpful against landslides and soil loss by preventing erosion.

Deep and narrow gullies are recommended to be put under permanent vegetation of grasses and trees. Bamboo plantation for productive and protective utilization of such degraded lands

is not a profitable option for local stake holders but also financially and economically viable policy option for funding agencies and Government and Non-Government agencies.

Program 3. Establishing Brushwood Check Dam for minimizing erosion hazard and reclamation of gullied land

Objectives

- a). To reduce the velocity of runoff.
- b) To prevent deepening and widening of the gully.
- c) To collect sedimentation and to recharge water table.

Expected outcomes

The proposed study is to Introducing of Brushwood Check Dam for minimizing erosion hazard and reclamation of gullied land will be conducted in a gully formed by the Southeast side of the Administrative Building of SCWMC, SRDI, Bandarban. It a large gully with uneven shaped. Larger gully rehabilitation requires more complex intervention to prevent continued erosion Check dams can be implemented in a stepped-approach for larger gully to slowly trap sedimentation and reclaim gully. In the hills of CHT, stone is not generally available everywhere, but brushes and unused trees are available Where stones are not readily available, Brushwood check dam can be constructed for slowly reclamation of the gullied land. Brushwood check dam increases absorption /infiltration of water into the soil. It also reduces the speed of runoff and therefore also reduces the erosive power of surface flows. Brushwood check dams allow for planting of crops once the dam is matured. It needs branches and plant materials/brushwood, ideally use of cuttings of trees that will strike fort the struts. Brushwood check dam can be build easily. But it needs for regular maintenance and repairing.

Program 4.STUDYING BROOM GRASS FOR CONTROLING SOIL EROSION AND ITS ECONOMIC VALUE AT CHT

Objectives

- a) To find out a significant source of income.
- b) To prevent frequent landslides, retain ground moisture and to increase fertility.
- c) To provide green forage for livestock.
- d) To increase ground water recharge and to enrich the ecological balance.

Justification

Thysanolaena maxima is a genus of plants in the grass family, the only genus in the tribe. It is locally known as Broom grass, Jharu ful, Fuljhru, Foruin etc. It's other names are Tiger Grass, Nepalese Broom Grass, Broom stick, Nepali amliso or kuchcho, jhadu (phool jhadu) in Hindi.

Broom grass is a significant source of income for subsistence communities, primarily for the women who collect it to manufacture and sell them as brooms. It can help the women to become more empowered by raising their financial status and lessening the burden of other tasks and thus it can ensure sincere impact of the ladies in the society. In addition to providing cash income when sold as brooms the plant provides a variety of uses to the farmers such as, the leaves provide green forage for livestock, the roots promote soil conservation, and the dried up stems can be used as stakes to support growing vegetables. Broom grass has a direct impact in preventing frequent landslides, helps to retain ground moisture and fertility, and can improve soil quality by reducing soil erosion. Broom grass grows in clumps and has many tangled up roots that grow to about one metre below the ground. This makes it highly effective in preventing soil erosion on hillsides as the grass is less likely to fall compared to other plants and trees that would have been planted there. The roots and leaves of the plant slow down water drops and the flow of water after heavy rain by absorbing the water in the soil Broom grass has the ability to crowd out invasive species when intercropped and is beneficial in retaining soil nutrients to re-grow vegetation. The grass also possesses numerous medicinal properties that are essential in subsistence communities. Broom grass does not compete for land with cereal crops so they can be grown simultaneously.

Growing broom grass on degraded land has been proven to help rehabilitate it as it helps retain ground moisture and promote fertility. There is no irrigation required to grow the grass and it does not produce any wastewater. No external inputs or energy is needed to grow the plant as it only requires human labour, which can be extensive in the first year of growing. Broom grass farming is highly recommended in new shifting cultivation systems on marginal lands to repair the damage from previous slash and burn methods. So it is very helpful to increase the local biodiversity in the communities and endangered animals can reappear.

Program 5. Studying effect of Natural vegetative strip (NVS) for minimizing soil erosion in cultivation of vegetables

Objectives:

1. To examine the effect of NVS on the maintenance of soil fertility and reducing soil erosion in moderate hill slope.
2. To examine the effect of NVS on crop productivity in hill slope.

Expected output/Justification

Minimization of soil erosion through natural vegetative strip (NVS) is an indigenous technology which used by the hill dwellers since time immemorial. The conservation of soil and water is essential for sustainable production, environment preservation and balanced eco system. Loss of soil by water erosion on sloping lands adversely affects the physical, chemical and biological properties of soils, leading to low crop productivity in this manner the experiment will established to control soil erosion and sustain crop productivity and aware the people as well as the peoples who involved to develop the people of the remote area.

Program 6. Development of Integrated Watershed Management at SCWMC in CHT (on going).

Program 7. Study on Management and economic value of *Schumannianthus dichotoma* (Murta /Patibet) in hilly Jhiri land at CHT (on going).

Program 8. Effect of different Hedge Species on Controlling Soil Erosion, Runoff, And Nutrient Mining Of Snack Gourd At Gentle Slope In CHT (on going).

Program 9. Studying Soil Loss and Yield Performance of Pineapple Based Jackfruit Orchard on Hill Slope following Contour line (on going).

4.2 Research activities of Salinity Management and Research Center (SMRC), Batiaghata, Khulna

Experiment 1. Selection of suitable sunflower varieties under dibbling cultivation in coastal saline soil

Abstract

The experiment was conducted at SMRC, Batiaghata, Khulna during rabi season of 2017-18 to identify suitable sunflower varieties by dibbling method for escaping high soil salinity in late drained land. Two sunflower varieties viz. BARI surjomukhi 2 and Suvarna were used in the experiment. Just after harvesting of T. Aman, sunflower seeds were sown in line by dibbling method. Balanced fertilizer was applied in the plots. Total amount of TSP, MOP, Gypsum and Zinc sulphate were applied 5 days before seed sowing as basal. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. The experiment was laid out in a Randomized Complete Block (RCB) design with three replications. The highest yield (1.55 t/ha) was obtained from BARI surjomukhi 2. The lowest yield was obtained from Suvarna (1.48 ton/ha) which was statistically similar with the variety BARI surjomukhi 2.

OBJECTIVES

- i) To identify suitable sunflower varieties by using dibbling method for escaping high soil salinity in late drained land.
- ii) To increase cropping intensity in coastal saline area by introducing sunflower in late drained land.

MATERIALS AND METHODS

The experiment was carried out at SMRC, Batiaghata, Khulna during Rabi season of 2017-2018. Two varieties of sunflower were used viz. BARI surjomukhi 2 and Suvarna. Just after harvesting of T. Aman, sunflower seeds were sown by dibbling method in line. Line to line and seed to seed distance was 60 cm and 30 cm respectively. Two Germinated seeds were dibbled in a hole. After 15 days of germination one healthy plant was kept and other one was removed. Sun flower seeds were sown on 30-11-2017 and after 40 days of emergence first weeding was done at 'Zo' condition by spade. Balanced fertilizer was used by considering initial soil status and requirements of sunflower. Total amount of TSP, MOP, Gypsum and Zinc sulphate were applied 5 days before seed sowing as basal. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. Weeding was also done when necessary. Insects and pests were controlled by using insecticides and pesticides when necessary. Sunflower was harvested on 06-04-2018.

Table 22. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				$\mu\text{g/g}^{-1}$			
7.9	2.23	0.21	0.140	21.15	23.18	0.91	0.58
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

Under dibbling condition, BARI surjomukhi 2 gave highest yield which was 1.55 t/ha. The lowest yield (1.48 t/ha) was obtained from variety Suvarna but the yield was also satisfactory.

Table 23. Yield of sunflower

Sunflower Varieties	yield (t/ha)
BARI surjomukhi 2	1.55
Suvarna	1.48
CV(%)	6.11

During experiment period soil salinity of experimental plot was increasing from sowing to harvesting. The highest soil salinity (EC: 8.94 dS/m) was recorded in March.

Table 24. Month wise soil salinity of Experimental plot.

Month	Soil Salinity: EC (dS/m)
November/2017	1.84
December/2017	2.36
January/2018	4.52
February/2018	6.07
March/2018	8.94

CONCLUSION

The experiment will be continued for the next year for confirmation of the result obtained in first year.

Experiment 2: Selection of suitable wheat varieties under zero tillage condition in coastal saline soil

Abstract

The experiment was carried out at SMRC, Batiaghata, Khulna during rabi season of 2017-18 to identify suitable wheat varieties under zero tillage condition for escaping high soil salinity in late drained land. The experiment was laid out in a Randomized Complete Block (RCB) design with three replications and two varieties. The wheat varieties were BARIGom 25 and BARIGom 26. Just after harvesting of T. Aman, the total amount of TSP, MOP, Gypsum and Zinc sulphate were applied as basal in wet condition. After 5 days wheat seeds were broadcast in the wet land under zero tillage condition. Urea was top dressed in three equal splits at 15, 40 and 60 days after seed sowing. The highest yield (2.64 t/ha) was obtained from BARI Gom 25. The lowest yield was obtained from BARI Gom 26 (2.31 ton/ha) which was statistically similar with that of BARIGom 25.

OBJECTIVES

- i) To identify suitable wheat varieties by zero tillage method for escaping high soil salinity in late drained land.
- ii) To increase cropping intensity in coastal saline area by introducing wheat in late drained land.

MATERIALS AND METHODS

The experiment was conducted at SMRC, Batiaghata, Khulna during Rabi season of 2017-2018. The varieties of wheat were BARI Gom 25 and BARI Gom 26. Just after harvesting of T. Aman, the total amount of TSP, MOP, Gypsum and Zinc sulphate were applied as basal in wet condition. After 5 days wheat seeds were broadcasted in the field. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. Thinning and first weeding was done after 20 days of sowing. Weeding was also done when necessary. Insects and pests were controlled by using insecticides and pesticides when necessary. Wheat was harvested on 30-03-2018.

Table 25. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				$\mu\text{g/g}^{-1}$			
7.7	2.27	0.27	0.132	20.17	29.98	0.98	0.29
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

Under zero tillage condition, highest yield was obtained from BARI Gom 25 (2.64 t/ha) and the lowest yield was obtained from BARI gom 26 (2.31 t/ha).

Table 26. Yield of wheat

Wheat varieties	yield (t/ha)
BARI Gom 25	2.64
BARI Gom 26	2.31
CV(%)	6.37

During experiment period soil salinity of experimental plot was observed to be increasing from sowing to harvesting. The highest soil salinity (EC: 9.14 dS/m) was recorded in March.

Table 27. Month wise soil salinity of Experimental plot.

Month	Soil Salinity: EC (dS/m)
November/2017	2.63
December/2017	4.18
January/2018	5.58
February/2018	7.22
March/2018	9.14

CONCLUSION

Under zero tillage condition, the growth and yield of wheat was satisfactory.

RECOMMENDATION

Under zero tillage condition, BARI Gom 25 and BARI Gom 26 may be recommended to farmer's field for cultivation in saline soil of coastal area.

Experiment 3: Effect of split application of DAP on the yield of kharif bottle gourd on raised ail in T. Aman rice field of saline area

Abstract

The experiment was conducted at SMRC, Batiaghata, Khulna during kharif season of 2017-18 to evaluate the effect of split application of DAP on the yield of kharif bottle gourd on raised ail in T. aman rice field of coastal saline soil. The experiment was designed with 4 treatments viz. T₀ : Control (Total P applied as basal), T₁ : 50% P as DAP solution applied 2 splits, T₂ : 50% P as DAP solution applied 4 splits and T₃ : 50% P as DAP solution applied 6 splits. The entire amount of TSP, MOP, Gypsum and Zinc sulphate were applied as basal for control (T₀). In case of Treatment T₁, T₂ and T₃, 50% dose of P was applied as basal and other 50% dose of P was applied as splits by DAP. The experiment was laid out in a Randomized Complete Block (RCB) design with three replications. The highest yield (51.59 t/ha) was obtained from T₁ treatment followed by T₂ (48.51 ton/ha) and T₃ (47.17 ton/ha). The lowest yield was obtained from T₀ treatment (42.62 ton/ha).

OBJECTIVES

- i) To evaluate the performance of split application of DAP solution in Kharif bottle gourd cultivated on raised ail of T. Aman rice
- iii) To introduce kharif bottle gourd in the coastal saline area.

MATERIALS AND METHODS

An experiment was carried out at SMRC, Batiaghata, Khulna during kharif-2 season, 2017. The experiment was laid out in RCBD design with three replications. There were four treatments in this experiment. The treatments were –T₀ : Control (Total P applied as basal), T₁ : 50% P as DAP solution applied 2 splits, T₂ : 50% P as DAP solution applied 4 split and T₃ : 50% P as DAP solution applied 6 split. Raised pits on ail in T. Aman field were made properly at 10-07-2017. Basal dose of manure and chemical fertilizers was applied in pit based on soil test value. Fertilizers were applied @ 130, 145, 30, 20, 15gm/pit of Urea, TSP, MOP, Gypsum and Zinc sulphate (heptahydrate), respectively. The entire amount of TSP, MOP, Gypsum and Zinc sulphate were applied as basal for control (T₀). In case of Treatment T₁, T₂ and T₃, 50% dose of P was applied as basal and other 50% dose of P was applied as splits by DAP. Urea was top dressed in three equal split at 15, 40 and 60 days after sowing. Pit to pit distance was 2.0 m. Germinated seeds were sown in pit at 21-08-2017. Intercultural operation was done as and when necessity. Insect and pest were controlled by using insecticides and pesticides. Green fruit collection was started from 15-10-2017 and ends at 12-12-2017.

Table 28. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				$\mu\text{g/g}^{-1}$			
8.1	1.12	0.29	0.101	3.21	86.97	0.53	0.82
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

In the experiment, the highest yield (51.59 t/ha) was obtained from the treatment T₁ where 50% of P was applied as DAP solution at 2 splits, followed by T₂ (48.51 t/ha) and T₃ (47.17 t/ha). The lowest yield was found from T₀ (42.62t/ha) treatment where full doses of Phosphorus was applied as basal.

Table 29. Yield of sweet gourd was under different treatments

Treatment	Bottle gourd yield (t/ha)
T ₀ (Control)	42.62c
T ₁ (50% P as DAP solution applied 2 split)	51.59a
T ₂ (50% P as DAP solution applied 4 split)	48.51b
T ₃ (50% P as DAP solution applied 6 split)	47.17b
CV(%)	8.37

CONCLUSION

The experiment may be continued for the next year for confirmation of the result.

Experiment 4: Effect of integrated nutrient management on soil salinity and yield of musk melon in saline soil

Abstract

The experiment was conducted at SMRC, Batiaghata, Khulna during kharif 1 season of 2017-18 to evaluate the effect of organic manure on soil salinity and yield of musk melon in saline soil. The variety of musk melon was local (Rongila). The experiment was designed with 7 treatments viz T₁: Control (STB chemical fertilizer), T₂ : Recommend cowdungwith chemical fertilizer (IPNS), T₃ : (Recommend + 25%) cowdungwith chemical fertilizer (IPNS), T₄ : (Recommend+50%)cowdungwith chemical fertilizer (IPNS), T₅ : Recommend poultry manurewith chemical fertilizer (IPNS), T₆ : (Recommend + 25%) polultry manurewith chemical fertilizer (IPNS), T₇ : (Recommend+ 50%) poultry manurewith chemical fertilizer (IPNS). The experiment was laid out in a Randomized Complete Block (RCB) design with three replications. The highest yield (26.05 t/ha) was obtained from T₇ treatment followed by T₄ (24.99 ton/ha) and T₆ (23.67 ton/ha). The lowest yield was obtained from T₁ treatment (18.43 ton/ha).

OBJECTIVES

- i) To identify effective organic manure with right dose to reduce soil salinity.
- ii) To find out the suitable combination of organic and inorganic fertilizer for optimizing the yield of musk melon.

MATERIALS AND METHODS

An experiment was conducted at SMRC, Batiaghata, Khulna. Musk melon was used as test crop. The experiment was laid out in RCB design with three replications having 6×4 m plot size. The land was prepared by 3 ploughing by power tiller. Pits were made properly. Manure and basal dose of fertilizer was applied during pit preparation. Pit to pit and line to line distance was 2.0 m. Germinated seeds (variety- deshi, Rangila) were sown in pit. Urea, TSP, MOP, Gypsum and Zinc sulphate were applied @ 110, 100, 15, 20, 20 gm/pit, respectively. The total amount of TSP, MOP, Gypsum and Zinc sulphate were applied as basal. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. Necessary intercultural operations were done as and when necessary. The ripe musk melon collection was started from 19-04-2017 and ended at 08-05-2017. After collection, the samples were analyzed statistically following MSTAT-c software.

Treatments: Seven treatments of the experiment were -

T₁ : STB chemical fertilizer

T₂ : Recommend cowdung with chemical fertilizer (IPNS)

T₃ : (Recommend + 25%) cowdung with chemical fertilizer (IPNS)

T₄ : (Recommend +50%) cowdung with chemical fertilizer (IPNS)

T₅ : Recommend poultry manure with chemical fertilizer (IPNS)

T₆ : (Recommend + 25%) poultry manure with chemical fertilizer (IPNS)

T₇ : (Recommend + 50%) poultry manure with chemical fertilizer (IPNS)

Table 30. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				μg/g ⁻¹			
7.5	1.54	0.26	0.089	6.46	147.80	0.36	0.72
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

Table 31. Nutrient status of different organic manure

Organic manure	pH	OC	Total N	P	K	S	Zn	B	C:N
		%							
Cowdung	7.5	19.2	2.0	1.3	0.8	0.03	0.05	0.15	9.6:1
Poultry manure	8.1	22.3	2.8	0.1	0.5	0.02	0.01	0.11	8.0:1

RESULTS AND DISCUSSIONS

It was observed that soil salinity was lowest in January and highest in April. Salinity data showed that organic manure contributed in decreasing soil salinity.

Table 32. Month wise soil salinity of the experimental plot during growing period

Treatments	Month wise Soil salinity(dS/m)				
	Jan/18	Feb/18	Mar/18	Apr/18	May/18
T ₁	4.7	7.9	11.8	16.6	10.9
T ₂	4.7	7.5	11.3	15.4	9.8
T ₃	4.7	7.2	11.1	15.2	9.5
T ₄	4.7	7.0	10.8	14.9	9.2
T ₅	4.7	7.4	11.2	15.3	9.6
T ₆	4.7	7.1	10.9	15.1	9.4
T ₇	4.7	6.9	10.7	14.8	9.0

From the experiment, highest yield (26.05 t/ha) was obtained from T₇ treatment followed by T₄ (24.99 ton/ha) and T₆ (23.67 ton/ha). The lowest yield was obtained from T₁ treatment (18.43 ton/ha).

Table 33. Yield of musk melon under different treatments

Treatment	Yield (t/ha)
T ₁ (STB chemical fertilizer)	18.43
T ₂ (Recommend cowdung) with chemical fertilizer(IPNS)	20.95
T ₃ (Recommend + 25%) cowdung with chemical fertilizer(IPNS)	22.82
T ₄ (Recommend +50%) cowdung with chemical fertilizer(IPNS)	24.99
T ₅ Recommend poultry manure with chemical fertilizer(IPNS)	21.80
T ₆ (Recommend + 25%) poultry manure with chemical fertilizer(IPNS)	23.67
T ₇ (Recommend + 50%) poultry manure with chemical fertilizer(IPNS)	26.05

CONCLUSION

Maximum Soil salinity was reduced by using poultry manure and plant growth & yield of musk melon was highest from the treatment T₇: (Recommend +50%) poultry manure with IPNS chemical fertilizer. But the yield obtained from T₄ was close to T₇.

RECOMMENDATION

For reducing soil salinity and getting better yield of musk melon farmers can use poultry manure or cowdung as organic manure.

Experiment 5. Selection of suitable kharif sweet gourd varieties in coastal saline soil

Abstract

The experiment was conducted at Salinity Management and Research Centre, Batiaghata, Khulna during kharif 2017-2018 to select suitable sweet gourd varieties namely Sweety (Laltirr Seed Limited), Sonia (Krisibid Seeds Limited), Super sweet (Northern Agriscience Limited), Lalima (Metal Agro Limited), Haland (Ispahani Agro Limited) and BARI Hybrid Mistikumra -1 for coastal saline soil. The highest yield (27.83 t/ha) was recorded from Haland variety followed by Sonia (24.80 t/ha), Sweety (23.13 t/ha), BARI hybrid misti kumra-1 (22.21 t/ha) and Lalima (11.97 t/ha) while Super sweet gave the lowest yield (7.69t/ha).

OBJECTIVES

- i) To select suitable sweet gourd varieties managing saline soil and water for coastal saline area.
- ii) To bring seasonally fallow land under crop production.

MATERIALS AND METHODS

An experiment was carried out at SMRC, Batiaghata, Khulna. . The experiment was laid out in RCBD design with three replications. The experiment was set up on kharif-1 season of 2018. Six hybrid varieties of sweet gourd namely Sweety (Laltirr Seed Limited), Sonia (Krisibid Seeds Limited), Super sweet (Northern Agriscience Limited), Lalima (Metal Agro Limited), Haland (Ispahani Agro Limited) and BARI Hybrid Mistikumra -1 cultivated for screening purpose. The land was prepared by three times ploughing by power tiller. Pits were made properly. Manure and basal dose of fertilizer was applied during pit preparation. Pit to pit and line to line distance was 2.0 m. Germinated seeds were sown in pit. Urea, TSP, MOP and Zinc sulphate (heptahydrate) were applied @ 111, 119, 23, 4 gm/pit, respectively. The total amount of TSP, MOP and Zinc sulphate were applied as basal. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. Necessary intercultural operations were done as and when necessary. Fruits (bitter gourd) collection was started from 21-05-2018 and ended at 26-05-2018. Data on yield components like fruit length, fruit diameter, individual fruit weight and no. of fruits per plant

were recorded from 10 randomly selected plants as per varieties and replications. After collection, the samples were analyzed statistically following MSTAT-c software.

Table 34. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				µg/g-1			
7.8	1.87	0.40	0.131	10.17	40.34	0.54	0.98
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

Among the six hybrid varieties of sweet gourd Haland gave statistical highest yield (27.83 t/ha) followed by Sonia (24.80 t/ha), Sweety (23.13 t/ha), BARI hybrid misti kumra-1 (22.21 t/ha) and Lalima (11.97 t/ha). The lowest yield was obtained from Super sweet (7.69t/ha).

Table 35. Yield of sweet gourd in saline soil

Varieties	Yield (t/ha)
Sweety	23.13b
Sonia	24.80b
Super sweet	7.69d
Lalima	11.97c
Haland	27.83a
BARI Hybrid Mistikumra -1	22.21b
CV (%)	6.07

Table 36. Month wise soil salinity in the experimental plot during cultivation

Year	Month wise Soil salinity (EC: dS/m)				
	Jan	Feb	Mar	April	May
2018	7.98	8.87	11.28	12.67	8.96

Soil salinity was increased from the month of January to April. The highest soil salinity was recorded in the month of April (12.67dS/m), followed by March (11.28dS/m), May (8.96dS/m) and February (8.87dS/m). The lowest soil salinity was observed in the month of January (7.98dS/m).

CONCLUTIONS AND RECOMMENDATIONS

Among six hybrid varieties of sweet gourd Haland, Sonia, Sweety and BARI hybrid mistikumra -1 may be recommended for cultivation in farmer's field under slightly to moderate soil salinity in coastal area.

Experiment 6. Selection of suitable kharif ribbed gourd varieties in coastal saline soil

Abstract

A field experiment was carried out at Salinity Management and Research Centre, Batiaghata, Khulna during kharif 2017-2018 cropping year to select suitable ribbed gourd varieties in coastal saline soil. Five hybrid varieties namely Hero (Laltirr Seed Limited), Sindabad (Getco Agro Vision Limited), Super moon (Northern Agri science Limited), Ishakha (Mollika Seed Company) and BARI Jhinga-02 were used in the study. Among the varieties highest yield (10.41 t/ha) was obtained from Hero followed by Sindabad (9.39 t/ha), BARI Jhinga-02 (6.78 t/ha) and Super moon (6.00 t/ha). The lowest yield was found from Ishakha variety which was 4.09 t/ha.

OBJECTIVES

- To select suitable rib gourd varieties managing saline soil and water for coastal saline area
- To bring seasonally fallow land under crop production.

MATERIALS AND METHODS

An experiment was carried out at SMRC, Batiaghata, Khulna during kharif-1 season, 2018. The experiment was laid out in RCBD design with three replications. Five hybrid varieties of rib gourd namely Hero (Laltirr Seed Limited), Sindabad (Getco Agro Vision Limited), Super moon (Northern Agri science Limited), Ishakha (Mollika Seed Company) and BARI Jhinga-02 were cultivated for selection of suitable variety in coastal saline soil. The land was prepared by three times ploughing by power tiller. Pits were made properly. Manure and basal dose of fertilizer was applied during pit preparation. Pit to pit and line to line distance was 2.0 m. Germinated seeds were sown in pit. Urea, TSP, MOP and Zinc sulphate (heptahydrate) were applied @ 77, 40, 10, 3 gm/pit, respectively. The total amount of TSP, MOP and Zinc sulphate were applied as basal. Urea was top dressed in three equal splits at 15, 30 and 55 days after sowing. Necessary intercultural operations were done as and when necessary. Fruits were collected after every 2 days interval. Crop harvest period was 01-04-2018 to 13-05-2018. Data on yield components like fruit length, fruit diameter, individual fruit weight and no. of fruits per plant were recorded from 10 randomly selected plants as per varieties and replications. After collection, the samples were analyzed statistically following MSTAT-c software.

Table 37. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				µg/g-1			
7.8	1.99	0.37	0.136	12.76	45.18	0.63	1.03
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

Soil salinity was increased from the month of January to April. The highest soil salinity was recorded in the month of April (11.22 dS/m), followed by March (9.96 dS/m), May (8.89 dS/m) and February (5.45 dS/m). The lowest soil salinity was observed in the month of January (4.18dS/m).

Table 38. Month wise soil salinity in experimental plot

Year	Month wise Soil salinity (EC: dS/m)				
	January	February	March	April	May
2018	4.18	5.45	9.96	11.22	8.89

From the experiment statistically highest yield (10.41 t/ha) was obtained from the variety Hero, followed by Sindabad (9.39 t/ha), BARI Jhinga-02 (6.78 t/ha) and Super moon (6.00 t/ha). The lowest yield was found from Ishakha (4.09 t/ha) which was statistically dissimilar with other varieties.

Table 39. Yield of ribbed gourd

Varieties	Yield (t/ha)
Hero	10.41a
Sindabad	9.39b
Super moon	6.00c
IshaKha	4.09d
BARI Jhinga-02	6.78c
CV(%)	5.41

CONCLUSIONS AND RECOMMENDATIONS

Among these five varieties of ribbed gourd, Hero and Sindabad may be recommended for cultivation in farmer's field under slightly to moderate soil salinity in coastal area.

Experiment 7. Selection of suitable kharif snake gourd varieties in coastal saline soil

Abstract

A field experiment was conducted at Salinity Management and Research Centre, Batiaghata, Khulna during kharif 2017-2018 cropping year to select suitable snake gourd varieties in coastal saline soil. There were five hybrid snake gourd varieties namely Asa (Rahim Afroz C I C Agro Limited), Surma (A R Malik Seeds pvt.Limited), Barnali(Mollika seed Company), Nagraj (Northern Agri science Limited) and BARIchichinga -1 were cultivated in this study. The highest yield (19.71 t/ha) was recorded from Asa variety followed by Barnali (19.56 t/ha), Surma (18.43 t/ha) and BARIchichinga -1(17.87 t/ha) while Nagrajgave the lowest yield (6.98t/ha).

OBJECTIVES

- To select suitable snake gourd varieties managing saline soil and water for coastal saline area.
- To bring seasonally fallow land under crop production.

MATERIALS AND METHODS

An experiment was carried out at SMRC, Batiaghata, Khulna. The experiment was laid out in RCBD design with three replications. The experiment was set up on kharif-1 season of 2018. Five hybrid varieties of snake gourd namely Asa (Rahim Afroz C I C Agro Limited), Surma (A R Malik Seeds pvt.Limited), Barnali (Mollika seed Company), Nagraj (Northern Agri science Limited) and BARI chichinga -1 were cultivated for screening purpose. The land was prepared by three times ploughing by power tiller. Pits were made properly. Manure and basal dose of fertilizer was applied during pit preparation. Pit to pit and line to line distance was 1.50 m. Germinated seeds were sown in pit. Urea, TSP, MOP and Zinc sulphate (heptahydrate) were applied @ 80, 65, 10, 5 gm/pit, respectively. The total amount of TSP, MOP and Zinc sulphate were applied as basal. Urea was top dressed in three equal splits at 15, 40 and 60 days after sowing. Necessary intercultural operations were done as and when necessary. Fruits collection was started from 06-05-2018 and ended at 14-06-2018. Data on yield components like fruit length, fruit diameter, individual fruit weight and no. of fruits per plant were recorded from 10 randomly selected plants as per varieties and replications. After collection, the samples were analyzed statistically following MSTAT-c software.

Table 40. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				µg/g-1			
7.9	1.87	0.37	0.129	7.85	49.86	0.59	0.84
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

Among the five varieties of snake gourd Asa gave the highest yield (19.71 t/ha) followed by Barnali (19.56 t/ha), Surma (18.43 t/ha) and BARI chichinga -1(17.87 t/ha). The lowest yield was obtained from Nagraj (6.98t/ha).

Table 41. Yield of bitter gourd in saline soil

Varieties	Yield (t/ha)
Hybrid (Asa)	19.71a
Hybrid (Surma)	18.43b
Hybrid (Barnali)	19.56a
Hybrid (Nagraj)	63.98c
BARI chichinga -1	17.87b
CV (%)	7.13

Table 42. Month wise soil salinity of experimental plot during cultivation

Year	Month wise Soil salinity (EC: dS/m)				
	Jan	Feb	Mar	April	May
2018	4.14	6.93	10.38	12.07	8.42

Soil salinity was increased from the month of January to April. The highest soil salinity was recorded in the month of April (12.07 dS/m), followed by March (10.38dS/m), May (8.42 dS/m) and February (6.93 dS/m). The lowest soil salinity was observed in the month of January (4.14dS/m).

CONCLUTIONS AND RECOMMENDATIONS

Among these four varieties of snake gourd like Asa (Rahim Afroz C I C Agro Limited), Barnali(Mollika seed Company), Surma (A R Malik Seeds pvt.Limited) and BARI chichinga -1 may be recommended for cultivation in farmer's field under slightly to moderate soil salinity in coastal area.

Experiment 8.Selection of suitable kharif bottle gourd varieties in coastal saline soil

Abstract

A field experiment was carried out at Salinity Management and Research Centre, Batiaghata, Khulna during kharif 2017-2018 cropping year to select suitable bottle gourd varieties in coastal saline soil. There were four bottle gourd varieties namely Hygreen (Metal Agro limited), Diana(Laltirr Seed Limited), Chisti (Getco Agro Vision Limited) and BARI Lau were cultivated in this study. Among the varieties statistically highest (97.36 t/ha) was obtained from Diana followed by Hygreen (94.77 t/ha) and Chisti (76.87 t/ha). The lowest yield was found from BARI Lau 4 variety which was 63.91 t/ha.

OBJECTIVES

- To select suitable bottle gourd varieties managing saline soil and water for coastal saline area
- To bring seasonally fallow land under crop production.

MATERIALS AND METHODS

An experiment was carried out at SMRC, Batiaghata, Khulna during kharif-2 season, 2018. The experiment was laid out in RCB design with three replications. Four hybrid varieties of cucumber (khira) namely Hygreen (Metal Agro limited), Diana (Laltirr Seed Limited), Chisti

(Getco Agro Vision Limited) and BARI Lau4 were cultivated for selection of suitable variety in coastal saline soil. The land was prepared by three times ploughing by power tiller. Pits were made properly. Manure and basal dose of fertilizer was applied during pit preparation. Pit to pit and line to line distance was 2.0 m. Germinated seeds were sown in pit. Urea, TSP, MOP and Zinc sulphate (heptahydrate) were applied @ 124, 154, 17, 4 gm/pit, respectively. The total amount of TSP, MOP and Zinc sulphate were applied as basal. Urea was top dressed in three equal splits at 15, 30 and 60 days after sowing. Necessary intercultural operations were done as and when necessary. Fruits were collected after every 3 days interval. Crop harvest period was 28-08-2017 to 22-11-2017. Data on yield components like fruit length, fruit diameter, individual fruit weight and no. of fruits per plant were recorded from 10 randomly selected plants as per varieties and replications. After collection, the samples were analyzed statistically following MSTAT-c software.

Table 43. Initial status of soil in experimental plot

pH	OM (%)	K meq/100 gm soil	Total N (%)	P	S	Zn	B
				µgg-l			
7.2	1.13	0.23	0.103	4.43	71.86	0.52	0.95
Critical level		0.12	0.12	7.00	10.00	0.60	0.20

RESULTS AND DISCUSSIONS

From the experiment statistically highest yield (97.36 t/ha) was obtained from the variety Diana, followed by Hygreen (94.77 t/ha) and Chisti (76.87 t/ha). The lowest yield was found from BARI Lau 4 (63.91 t/ha) which was statistically dissimilar with other varieties.

Table 44. Yield of sweet gourd under different varieties

Varieties	Yield (t/ha)
Hygreen	94.77a
Diana	97.36a
Chisti	76.87b
BARI Lau4	63.91c
CV (%)	5.89

During experiment period soil salinity of experimental plot was non saline condition due to rainy season.

Table 45. Month wise soil salinity in experimental plot

Year	Month wise Soil salinity (EC: dS/m)					
	June	July	Aug	Sep	Oct	Nov
2017	2.1	1.8	1.7	1.7	1.9	2.2

CONCLUSIONS AND RECOMMENDATIONS

Among the four varieties of bottle gourd, Diana and Hygreen may be recommended for cultivation in farmers field in coastal area.



Experiments with salt tolerant varieties of vegetables and field crops at SMRC, Batiaghata, Khulna

Proposed Research program (2018-19)

Expt. 1: Study on the performance of spices and vegetables under multi-layer crop production in high land of coastal saline area

Expt. 2: Study on the performance of brinjal under different ratio of manure and soil in flying bed of coastal saline area

Expt. 3: Study on the performance of tomato under different ratio of organic manure in flying bed of coastal saline field

Expt. 4: Study on the performance of chilli under different ratio of manure and soil in flying bed of coastal saline area

Expt. 5: Study on the performance of yard long bean under different ratio of manure and soil in flying bed of coastal saline area

Expt. 6: Study on the performance of yard long bean under organic agricultural system in coastal saline area

Expt. 7: Study on the performance of bitter gourd under organic agricultural system in coastal saline area

Expt. 8: Effect of integrated nutrient management on bitter gourd in plastic container

Expt. 9: Study on salt accumulation in soil by using different strength of saline water

Expt. 10: Screening of some cucumber varieties

Chapter 5: Activities of Regional Laboratories

Program 1. Analysis of Soil, Water, Plant & Fertilizer Samples

Introduction

Productivity of limited land and soil resources of Bangladesh is gradually declining because of indiscriminate use of chemical fertilizers, inadequate soil and water management and overall irrational use of lands. Every year a considerable portion of agricultural land is being converted to non-agricultural land due to unplanned infrastructure developments. To ensure food security of ever growing population of Bangladesh only option left is to increase crop production per unit area per unit time. This can be attained by selecting crops/cropping pattern on the basis of land and soil qualities, agro-climate, and use of balanced fertilizer (both chemical & organic) on soil test and demand basis, ensuring proper crop management practices.

In the Regional Laboratories soil, water and plant samples were analyzed to fulfill the following objectives.

Objectives:

- i) To recommend location specific balanced fertilizer doses for farmers and other stakeholders based on soil test result and crop demand.
- ii) To provide analytical support in SRDI's regular Nirdeshika Updating program, soil and water salinity monitoring program.

Determination of different elements

The soil samples received were analyzed following standard methods to determine pH, EC, organic carbon, exchangeable acidity, texture, total N, available P, K, S, Mg, Zn, Cu, Fe, Mn & B. Fertility status of soil is calculated from analytical data.

Plant materials were analyzed for determining their total N, P, K, Ca, Mg, Na, Fe, Mn, Cu & B content following standard methods. Nutrient deficiencies/toxicities may be inferred from the analytical data.

Water samples are analyzed for determining their irrigation suitability.

Fertilizer samples are analyzed for determining their ingredients as specified by the manufacturers. This is important for quality control of fertilizers.

Achievements in the year 2017-18

Dhaka, Comilla, Rajshahi and Khulna laboratories analyzed 6,415 nos. of soil samples and 62,353 nos. ingredients; 259 nos. of water sample and 1,077 ingredients; 5 plant samples and 25 ingredients; 2,239 nos. of fertilizer samples and 5,168 ingredients (Table-62). During this period earning non-tax revenue amounting to Tk.11,59,485.00 as analysis fee (Table-62).

Other static laboratories located at Barisal, Dinajpur, Bogra, Jamalpur, Noakhali, Kushtia, Faridpur, Jhenaidah, Mymensingh, Sylhet and Chittagong analyzed 11,677 nos. of soil samples and 81,558 nos. of ingredients (Table-65). During this period 6,503 nos. of Fertilizer Recommendation Cards distributed (Table-66) and non-tax revenue amounting Tk.4,25,639.00 was earned (Table-67).

Table 46. Number of soil samples & ingredients analyzed in 4 Regional Laboratories

Sl No.	Program/source of Samples	Target				Number of soil samples & ingredient analyzed							
		Dhaka	Comilla	Rajshahi	Khulna	Dhaka Regional laboratory		Comilla Regional laboratory		Rajshahi Regional laboratory		Khulna Regional laboratory	
						Samples	Ingredients	Samples	Ingredients	Samples	Ingredients	Samples	Ingredients
1.	Farmer's soil	600	800	500	600	30	239	164	954	1,040	7,397	159	1,405
2.	SRDI's program												
	a) Upazilla Nirdehaka	1000	800	1000	750	836	8,975	1,038	14,532	1,087	10,911	622	8,046
	b) Monitoring	200	50	104	80								
	c) Block survey	-	-	-	-								
	d) FRC	100	50	200	-								
	e) Quality control	100	100	108	70	93	1,105	173	1,748	24	96	50	236
	f) Other program	-	200	-	-	104	658			32	416	178	527
3.	Other organization	200	300	100	200	292	2,212	633	2,877	142	1,294	59	434
4.	SFFM	-	-	-	-								
	Total	2,200	2,300	3,012	1,700	1,355	13,189	1,667	18,402	2,325	20,114	1,068	10,648

Note: Other Organization include GO's, NGO's, NARS organizations, educational institutes etc.

Table 47. Number of Water samples & ingredients analyzed in 4 Regional Laboratories

Sl No.	Program/source of Samples	Target				Number of water samples & ingredient analyzed							
		Dhaka	Comilla	Rajshahi	Khulna	Dhaka Regional laboratory		Comilla Regional laboratory		Rajshahi Regional laboratory		Khulna Regional laboratory	
						Samples	Ingredients	Samples	Ingredients	Samples	Ingredients	Samples	Ingredients
1.	Farmer's											204	869
2.	Different organization	100	50	100	40	0	0	0	0	55	208	-	-
	Total	100	50	100	40	0	0	0	0	55	208	204	869

Note: Other Organization include GO's, NGO's, NARS organizations, educational institutes etc.

Table 48. Number of fertilizer samples & ingredient analyzed in 4 Regional Laboratories

Sl No.	Program/source of Samples	Target				Number of water samples & ingredient analyzed							
		Dhaka	Comilla	Rajshahi	Khulna	Dhaka Regional laboratory		Comilla Regional laboratory		Rajshahi Regional laboratory		Khulna Regional laboratory	
						Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients
2.	Different organization	600	400	800	500	429	1308	775	1543	695	1163	340	1154
Total		600	400	800	500	429	1308	775	1543	695	1163	340	1154

Note: Other Organization include GO's, NGO's, NARS organizations, educational institutes etc.

Table 49. Number of plant samples & ingredient analyzed in 4 Regional Laboratories

Sl No.	Program/source of Samples	Target				Number of plant samples & ingredient analyzed							
		Dhaka	Comilla	Rajshahi	Khulna	Dhaka Regional laboratory		Comilla Regional laboratory		Rajshahi Regional laboratory		Khulna Regional laboratory	
						Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients
2.	Different organization	100	50	100	140	0	0	0	0	4	24	1	1
Total		100	50	100	140	0	0	0	0	4	24	1	1

Note: Other Organization include GO's, NGO's, NARS organizations, educational institutes etc.

Table 50. Number of soil samples & ingredients analyzed in 4 regional laboratories

Sl No	Program/source of Samples	Number of soil samples & ingredient analyzed								Total		Revenue Earning (BDT)
		Dhaka Regional laboratory		Comilla Regional laboratory		Rajshahi Regional laboratory		Khulna Regional laboratory		Samples	Ingre-dients	
		Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients	Samples	Ingre-dients			
Soil samples	Other organization	1,355	13,189	1,667	18,402	2,325	20,114	1,068	10,648	6,415	62,353	11,59,485.00
Plant	Other organization	0	0	0	0	4	24	1	1	5	25	
Water	Other organization	0	0	0	0	55	208	204	869	259	1,077	
Fertilizer	Other organization	429	1308	775	1543	695	1163	340	1154	2,239	5,168	
Total		1,784	14,497	2,442	19,945	3,079	21,509	1,613	12,672	8,918	68,623	

Note: Other Organization include GO's, NGO's, NARS organizations, educational institutes etc.

Table 51. Number of soil samples and ingredients analyzed in different static laboratories

Sl. No.	Name of the Laboratory	No. of soil sample analyzed	No. of ingredient analyzed
1.	Regional Laboratory Barisal	1,166	7,437
2.	Regional Laboratory Bogra	1,448	8,139
3.	Regional Laboratory Chittagong	1,066	5,380
4.	Regional Laboratory Dinajpur	1,984	20,583
5.	Regional Laboratory Faridpur	882	3,568
6.	Regional Laboratory Jamalpur	553	2,212
7.	Regional Laboratory Jhenaidah	972	9,024
8.	Regional Laboratory Kushtia	806	4,198
9.	Regional Laboratory Mymensingh	569	3,504
10.	Regional Laboratory Noakhali	1,056	7,956
11.	Regional Laboratory Sylhet	1,175	9,557
Total		11,677	81,558

Table 56. Number of FRC prepared and distributed among the farmers

Sl. No.	Name of the Laboratory	No. of FRC prepared and distributed
1.	Regional Laboratory Barisal	514
2.	Regional Laboratory Bogra	783
3.	Regional Laboratory Chittagong	916
4.	Regional Laboratory Dinajpur	747
5.	Regional Laboratory Faridpur	841
6.	Regional Laboratory Jamalpur	502
7.	Regional Laboratory Jhenaidah	291
8.	Regional Laboratory Kushtia	432
9.	Regional Laboratory Mymensingh	568
10.	Regional Laboratory Noakhali	360
11.	Regional Laboratory Sylhet	549
Total		6,503

Table 57. Non-tax revenue earned as soil analysis fee

Sl. No.	Name of the Laboratory	Non-tax revenue Income (BDT)
1.	Regional Laboratory Barisal	37,957.00
2.	Regional Laboratory Bogra	59,164.00
3.	Regional Laboratory Chittagong	41,187.00
4.	Regional Laboratory Dinajpur	49,486.00
5.	Regional Laboratory Faridpur	14,700.00
6.	Regional Laboratory Jamalpur	12,075.00
7.	Regional Laboratory Jhenaidah	16,569.00
8.	Regional Laboratory Kushtia	10,130.00
9.	Regional Laboratory Mymensingh	29,060.00
10.	Regional Laboratory Noakhali	9,571.00
11.	Regional Laboratory Sylhet	9,557.00
Total		4,25,639.00

Chapter 6: Project Activities

6.1. Soil Research and Strengthening of Soil Research Facilities (SRSRF) Project

Project information

Implementing organization	: Soil Resource Development Institute (SRDI)
Administrative Ministry	: Ministry of agriculture
Name of Project	: Soil Research and Strengthening of Research Facilities (SRSRF)
Duration	: January, 2018- December, 2022
Approval date of ECNEC	: 02.01.2018
Administrative approval date	: 08.04.2018
Estimated Budget of Project	: 6308.25 Lac BDT
Economic Code	: 5-4345-5000
ADP allocation for 2018-19	: 1420.00 Lac BDT

Objectives of the Project:

1. To generate soil and land related data base for agricultural development plans & schemes through the research on problem soil and prepare of Union Sahayika.
2. To conduct research on problem soil management. Specially research on-
 - a. Hill Soil conservation and management techniques.
 - b. Technology development and extend on saline soil management for the sustainable agriculture.
 - c. Acidic Soil management through liming and cultural practices for sustainable agricultural development.
3. To implement adaptive research trial in the farmers field using the balanced fertilizer according to Upazila Nirdeshika & Union Sahayika.
4. Modernization of SRDI laboratories.
5. To established wave based virtual soil museum.
6. To arrange training for up-scaling capacity of farmers and extension people for soil management practices and awareness building through motivational approach.

During 2017-2018 fiscal year BDT 134.00 lac was allocated while expenditure was 133.42 (99.56%).

Achievement during 2017-2018

Sl. No.	Item	Achievement
Research		
01.	Research on Hill Soil conservation and management	5 research on going
02.	Research on Saline Soil management	4 research on going
03.	Site selection for acid soil management research	16 upazila of 8 district

Sl. No.	Item	Achievement
Printing Report		
01.	Printing of “Land, Soil and Fertilizer Recommendation guide” (Union Sahayika)	50 Union
Training		
01.	One day training on Soil samples collection and balanced fertilizer use for farmer, UDC entrepreneur & NGO worker	1900 (76 batch)
02.	One day training on adulterated fertilizer identification for Fertilizer dealer, SAAO, & NGO worker	250 (10 batch)
Seminar/Workshop/Conference		
01.	Group discussion	1
02.	Inception workshop	1
Modernization of Lab, maintenance of equipments and procurement of chemicals		
01.	Purchase of lab equipments	17 units of 4 items
02.	Maintenance of equipments	6 items
03.	Procurement of chemicals	51 kg various chemical 30 liter various acid
Procurement of Office furniture and computer equipments		
01.	Laptop/ Notebook	4
02.	Color printer	1
03.	Multimedia projector	1
04.	Photocopier	1
05.	Digital Camera	1

Program for 2018-2019

Sl. No.	Item	Target
Research		
01.	Research on Hill Soil conservation and management	17 research
02.	Research on Saline Soil management	7 research
03.	Research on acid soil management	32 research
Printing Report		
01.	Printing of “Land, Soil and Fertilizer Recommendation guide” (Union Sahayika)	84 Union
02.	Fertilizer recommendation festoon	84 Union

Training		
01.	One day training on Soil samples collection and balanced fertilizer use for farmer, UDC entrepreneur & NGO worker	1900 (76 batch)
02.	One day training on adulterated fertilizer identification for Fertilizer dealer, SAAO, & NGO worker	250 (10 batch)
03.	TOT-1: Six days training on Use of Upazila Nirdeshika for DAE officer	100 (4 batch)
03.	TOT-2: Three days training on Use of Upazila Nirdeshika for DAE officer	50 (2 batch)
04.	Six days training on Use of Upazila Nirdeshika for SAAO	630 (21 batch)
05.	5 days training on Soil conservation and hill Soil management in Nepal	4 SRDI Scientist
06.	Study tour in abroad	6 personnel
Seminar/Workshop/Conference		
01.	Group discussion	1
02.	Review workshop	1
03.	Attend workshop in abroad	1
Advertisement in print and Electronic media		
01	Leaflet/booklet/poster/folder/signboard for mass awareness	continuing
02.	Short film documentary on Soil test, use of balanced fertilizer and adulterated fertilizer	continuing
Modernization of Lab, maintenance of equipments and procurement of chemicals		
01.	Purchase of lab equipments	114 units of 24 items
02.	Maintenance of equipments	Need basis
03.	Procurement of chemicals	Need basis
Procurement of Office equipments, furniture and computer equipments		
01.	Computer, printer, GIS accessories	5
02.	LED TV	1
03.	Fax machine	1
04.	Water purifier	1
05.	Furniture	8 items

6.2 Development of Upazila Land Suitability Assessment and Crop Zoning System of Bangladesh (SRDI-part)

A. Summary of the Project

The objective of the project is to provide crop suitability information to different stakeholders including farmers on the choice of rotation of their crops that ensures the increase of food production and maximizes the farmer's income. More specifically the project objectives are:

- a) Creating, updating and validation of the land/crop suitability databases in order to derive appropriate farming practices for sustainable socio-economic condition.
- b) Development of an online GIS based software for cropland suitability assessment.

Crop suitability assessment primarily based on land and soil data-information, which may be termed as edaphic suitability. Ultimately crop suitability analysis of an area for a specific crop or cropping pattern is the combination of edaphic suitability, agro-climatic suitability and economic viability.

The Soil Resource Development Institute (SRDI) as a generator of soil and land information and as a partner of this project will be responsible for providing updated soil and land information data. For this SRDI will carry out survey for updating soil and land, nutrient status and other relevant data of Upazila Nirdeshika of the selected Upazilas. Further, they will be involved for collection of soil samples and analysis /interpretation of soil and land characteristics, soil profile information including fertilizer requirement of different crops and cropping patterns as per soil fertility status. The updated data as generated will be incorporated into the Upazila land resources data base which would be used for crop/land suitability assessment analysis.

More specifically activities of SRDI includes, semi-detailed soil survey, collection of soil samples, analysis of samples in the laboratories, map preparation and finally preparation of report known as “Soil and Land Resources Utilization Guide” popularly known as “Upazila Nirdeshika”. Land and soil resource database will use in analyzing location specific edaphic suitability of a crop or cropping pattern. Chemical data of soil samples will be used for location specific fertilizer recommendations for suitable crops and cropping patterns.

Moreover, SRDI will be involved in ground truthing or field validation. Ground truthing will be conducted on the crop suitability of the major crops grown in the study areas. The parameters, which are responsible for rating, will be checked in the field.

Soil Resource Development Institute will conduct semi-detailed soil survey in 60 upazilas in three years of project tenure (20+20+20) in order to update land and soil resources database. Every year they will conduct soil survey in 20 Upazilas under project fund and survey will conduct in another 30 upazilas under SRDI’s revenue budget.

SRDI will provide updated land and soil resources database including maps and reports of 300 upazilas for this project out of this SRDI already updated database of 150 upazilas before inception of this project.

B. Basic Project Information:

- i. Project Title -“**Upazila Land Suitability Assessment and Crop Zoning System of Bangladesh (SRDI-part)**”
- ii. Project Code/ID Number-- CRP-V
- iii. Name of Principal Investigator: Bidhan Kumar Bhandar
- iv. Name of Co Investigator: Mohammad Moniruzzaman
- v. Name of the applying organization with address- Soil Resource Development Institute (SRDI),Krishi Khamar Sarak, Dhaka-1215.
- vi. Project duration (year/months) From June, 2017 to May,2020
- vii. Project commencement date (As per MOU) - June,2017
- viii. Project Locations/Sites-All over Bangladesh

C. Specific Project Objective(s): (As per FRPP/PIR)

The objective of the project is to provide crop suitability information to different stakeholders including farmers on the choice of rotation of their crops that ensures the increase of food production and maximizes the farmer’s income. More specifically the project objectives are:

- a) Creating, updating and validation of the land/crop suitability databases in order to derive appropriate farming practices for sustainable socio-economic condition.
- b) Development of an online GIS based software for cropland suitability assessment.

D. Research Approaches and Methodologies:

i) Approaches:

Existing land type map will be updated using DEM data collected from the primary and/or secondary sources with high precision using GIS technology. SRDI will take part in ground truthing work to validate the land type boundary. Socio economically suitable cropping patterns generated by the developed crop zoning software will be validated in the field where SRDI will work as a member of the validation team.

A GIS based crop suitability assessment software and the associated tools will be developed as online application. The updated crop suitability database will be utilized for preparation of land suitability maps of the major cropping patterns of the selected Upazilas. Validation of the maps will be carried out in the fields where SRDI will contribute with its resource and expertise.

ii) Methodologies:

One of the important activities of the project is ground truthing or field validation. Ground truthing will be conducted on the crop suitability of the major crops grown in the study areas. The parameters, which are responsible for rating, will be checked in the field. In this respect the primary database for this area will be helpful. Also, inputs from knowledgeable farmers will help for carrying out this work. Socio-economic parameters and economically best suitable cropping patterns generated by the software will also be validated in the field. SRDI will also contribute with its knowledge and expertise in this process. However, BARC will plan and coordinate this activity along with other project partners.

Main responsibility of SRDI is to update soil and land resources database and also nutrient database through semi-detailed soil survey and provide updated database to BARC for crop suitability analysis. 60 upazila have been selected for database updating and updating survey. Each year 20 Upazilas will be surveyed for updating soil and land resource data for the project using project fund. SRDI will interpret updated database of all 300 upazilas selected under this project and process it in an appropriate form for crop zoning project. All the edaphic characteristics will be generated through SRDI. Each year updated database of 100 upazilas will be processed and provide it to Crop Zoning Project of BARC for crop suitability analysis.

